



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/28, 9/107	A1	(11) International Publication Number: WO 99/40932
		(43) International Publication Date: 19 August 1999 (19.08.99)

(21) International Application Number: PCT/CA99/00106

(22) International Filing Date: 5 February 1999 (05.02.99)

(30) Priority Data:

09/021,114	10 February 1998 (10.02.98)	US
09/216,733	21 December 1998 (21.12.98)	US

(71)(72) Applicant and Inventor: MODI, Pankaj [CA/CA]; 519
Golf Links Road, Ancaster, Ontario L9G 4X6 (CA).(74) Agent: BARRIGAR & MOSS; Suite 901, 2 Robert Speck
Parkway, Mississauga, Ontario L4Z 1H8 (CA).(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR,
BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE,
GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW,
MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO
patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR,
IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published

*With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: MIXED MICELLAR PHARMACEUTICAL DELIVERY SYSTEM AND METHOD OF PREPARATION

(57) Abstract

A mixed micellar pharmaceutical formulation includes a micellar proteinic pharmaceutical agent, an alkali metal C8 to C22 alkyl sulphate, alkali metal salicylate, a pharmaceutically acceptable edetate and at least one absorption enhancing compounds. The absorption enhancing compounds are selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof. The amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

*(Referred to in PCT Gazette No. 13/2000, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	NI	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

MIXED MICELLAR PHARMACEUTICAL DELIVERY SYSTEM AND METHOD OF PREPARATION

Field of the Invention

The present invention relates to an improved
5 delivery system for the administration of large-molecule
pharmaceuticals, e.g. peptidic drugs, vaccines and
hormones. In particular it relates to pharmaceuticals
which may be administered through the oral and nasal
membranes.

10 Background to the Invention

In spite of significant efforts in academic and
commercial laboratories, major breakthroughs in oral
peptide and protein formulation have not been achieved.
Relatively little progress has been made in reaching the
15 target of safe and effective oral formulations for
peptides and proteins. The major barriers to developing
oral formulations for proteins and peptides include poor
intrinsic permeability, lumenal and cellular enzymatic
degradation, rapid clearance, and chemical stability in
20 the gastrointestinal (GI) tract. Pharmaceutical
approaches to address these barriers, which have been
successful with traditional small, organic drug
molecules, have not readily translated into effective
peptide and protein formulations. Although the
25 challenges are significant, the potential therapeutic
benefits remain high especially in the field of diabetes
treatment using insulin.

Scientists have explored various administration
routes other than the injection for proteins and
30 peptides. These routes include oral, intranasal, rectal,
vaginal cavities for the effective delivery of large
molecules. Out of the above four mentioned routes oral

and nasal cavities have been of greatest interest to scientists. Both the oral and nasal membranes offer advantages over other routes of administration. For example, drugs administered through these membranes have
5 a rapid onset of action, provide therapeutic plasma levels, avoid first pass effect of hepatic metabolism, and avoid exposure of the drug to hostile GI environment. Additional advantages include easy access to the membrane sites so that the drug can be applied,
10 localized and removed easily. Further, there is a good potential for prolonged delivery of large molecules through these membranes.

The oral routes have received far more attention than has the other routes. The sublingual mucosa
15 includes the membrane of ventral surface of the tongue and the floor of the mouth whereas the buccal mucosa constitutes the lining of the cheek. The sublingual mucosa is relatively permeable thus giving rapid absorption and acceptable bioavailability of many drugs.
20 Further, the sublingual mucosa is convenient, acceptable and easily accessible. This route has been investigated clinically for the delivery of a substantial number of drugs.

The ability of molecules to permeate through the
25 oral mucosa appears to be related to molecular size, lipid solubility and peptide protein ionization. Small molecules, less than 1000 daltons appear to cross mucosa rapidly. As molecular size increases, the permeability decreases rapidly. Lipid soluble compounds are more
30 permeable than non-lipid soluble molecules. Maximum absorption occurs when molecules are un-ionized or

neutral in electrical charges. Therefore charged molecules present the biggest challenges to absorption through the oral mucosae.

Most proteinic drug molecules are extremely large
5 molecules with molecular weight exceeding 6000 daltons. These large molecules have very poor lipid solubility and are practically impermeable. Substances that facilitate the absorption or transport of large molecules (>2000 daltons) across biological membranes
10 are known as the enhancers, (Lee et al., Critical Reviews in Therapeutic drug Carrier Systems, 8, 91, 1991; Lee et al., Critical Reviews in Therapeutic drug Carrier Systems, 8, 115, 1991, 1992). Enhancers may be characterized as chelators, bile salts, fatty acids,
15 synthetic hydrophillic and hydrophobic compounds, and biodegradable polymeric compounds.

Various mechanisms of action of enhancers have been proposed. These mechanisms of action, at least for protein and peptidic drugs include (1) reducing
20 viscosity and/or elasticity of mucous layer, (2) facilitating transcellular transport by increasing the fluidity of the lipid bilayer of membranes, and (3) increasing the thermodynamic activity of drugs (Critical Rev, 117-125, 1991, 1992).

25 Many enhancers have been tested so far and some have found to be effective in facilitating mucosal administration of large molecule drugs. However, hardly any penetration enhancing products have reached the market place. Reasons for this include lack of a
30 satisfactory safety profile respecting irritation, lowering of the barrier function, and impairment of the

mucocilliary clearance protective mechanism. The main factor to be considered in the use of enhancers especially related to bile salts, and some protein solubilizing agents is extremely bitter and unpleasant taste. This makes their use almost impossible for human consumption on a daily basis. Several approaches were utilized to improve the taste of the bile salts based delivery systems, but none one of them are commercially acceptable for human consumption to date. Among the approaches utilized includes patch for buccal mucosa, bilayer tablets, controlled release tablets, use of protease inhibitors, buccally administered film patch devices, and various polymer matrices.

The basic problem associated with the above technologies is the use of large quantities of bile acids and their salts to promote the transport of the large molecules through membranes in the form of localized delivery system using patches or tablets. In spite of using protease inhibitors and polymer coatings the technologies failed to deliver proteinic drugs in the required therapeutic concentrations. Further, the problem is compounded because of the localized site effect of the patch which resulted in severe tissue damage in the mouth. Most attempts were made to deliver large molecules via the oral, nasal, rectal, and vaginal routes using single bile acids or enhancing agents in combination with protease inhibitors and biodegradable polymeric materials. However, it is extremely difficult to achieve therapeutic levels of proteinic drugs using these formulations. As single enhancing agents fails to loosen tight cellular junctions in the oral, nasal,

rectal and vaginal cavities for a required period of time to allow passage of large molecules through the mucosal membranes without further degradation. This problem makes it impractical to use the above mentioned systems for a commercial purpose.

In order to overcome the above mentioned problem of the bitter taste, irritation and the penetration of large molecules through the sublingual, buccal and GI tract mucosal lining, a system has now been designed where protein drug was encapsulated in mixed micelles made up of combination of enhancers, e.g. yolk proteins (lecithins). This system allows opening of the paracellular junctions (tight junctions) in oral as well as in GI tract by GI motility movement with high degree of protease activity preserved and protecting molecules from premature degradation in the hostile acidic and proteolytic GI environment.

It is believed that the mixed micelles encapsulate molecules with high degree of efficiency (>90% encapsulation). These mixed micelles are extremely small in the size (1 nm to 10 nm), and are smaller than the pores of the membranes in the oral cavity or the GI tract. It is therefore believed that the extremely small size of mixed micelles helps encapsulated molecules penetrate efficiently through the mucosal membranes of the oral cavity.

The absorption of proteins and peptides is believed to be enhanced by the diffusion of large molecules entrapped in the mixed micellar form through the aqueous pores and the cell structure perturbation of the tight paracellular junctions.

The amount of physiologically peptide or protein in the compositions of this invention is typically a quantity that provides an effective amount of the drug to produce the physiological activity (therapeutic plasma level) for which peptide or protein is being administered. In consideration of the fact that the bioavailability of any active substance can never be 100%, that is to say the administered dose of the active drug is not completely absorbed, it is preferable to incorporate slightly larger amount than the desired dosage. Where the dosage form is a spray (aerosol) or the like which is repeatedly dispensed from the same container, it is recommendably so arranged that the unit dose will be slightly greater than the desired dose. It should be understood that dosage should vary with species of warm blood animals such as man, domestic animals, and their body weights. Although the composition of this invention is prepared as the microfine droplets (1 to 10 nm or less) by the virtue of its preparation methods used and suitable combinations of enhancer compound characteristics. The utilization of atomizer or aerosol spray devices (metered dose inhalers or nebulizers) may be useful to further a sufficient reduction of particle size for effective inhalation from the nasal or oral cavity so the drug may successfully absorbed or reach to the specific site.

The therapeutic composition of the present invention can be stored at room temperature or at cold temperature. Storage of proteinic drugs is preferable at the cold temperature to prevent the degradation of the drugs and to extend their shelf life. While the mixed

micellar therapeutic composition of the invention is applied to the mucosal membranes, the sites of administration may be the same as those used for the usual mucosal therapeutic preparation. Generally, oral, 5 transdermal and nasal are the favourite sites of the administration but the composition can be applied to the rectal and vaginal mucosa. According to the physiologically active peptide or protein used, the dosage form and the site of administration, a specific 10 administration method can be selected.

As used herein, the term "edetate" is used herein to refer to pharmaceutically acceptable salts of ethylenediaminetetraacetic acid.

It has also been found that improvements in 15 penetration and absorption of mixed micellar formulations can be achieved by mixing the mixed micellar formulation with propellants such as tetrafluoroethane, heptafluoroethane, dimethylfluoropropane, tetrafluoropropane, butane, 20 isobutane, dimethyl ether and other non-CFC and CFC propellants. Preferably they are delivered through metered dose spray devices. Metered dose inhalers are known and are a popular pulmonary drug delivery form for some drugs. The present formulation, including the 25 propellant is intended to improve the quality of absorption, stability and performance of many formulations. The compositions have been selected to give enhancement in the penetration through pores, and facilitate absorption of the drugs to reach therapeutic 30 levels in the plasma. The present formulation may be absorbed buccally, by ensuring that the person does not

inhale the formulation as it is sprayed. One of the other benefits of using an atomizer or inhaler is that the potential for contamination is minimized because the devices are self contained.

5 Summary of the Invention

Accordingly the present invention provides a mixed micellar pharmaceutical formulation, having a pH of between 6.0 and 7.0 comprising a proteinic pharmaceutical agent in micellar form, water, an alkali metal lauryl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one micelle forming compound selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

In an embodiment, the alkali metal lauryl sulphate, the edetate and the alkali metal salicylate are each in a concentration of from 2 to 5 wt./wt.% of the total

formulation.

In one embodiment, the edetate is an alkali metal edetate. Preferably the alkali metal edetate is be selected from the group consisting of disodium edetate, 5 dipotassium edetate, and combinations thereof.

In another embodiment, the alkali metal lauryl sulphate is sodium lauryl sulphate.

In a further embodiment, the alkali metal salicylate is sodium salicylate.

10 In another embodiment, the lecithin is selected from the group consisting of saturated phospholipid, e.g. Phospholipon-H (trade mark) saturated phospholipid, unsaturated phospholipid, e.g. Phospholipon-G (trade mark) unsaturated phospholipid, phosphatidylcholine, 15 phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, and lysolecithin.

In one embodiment, one of the absorption enhancing compounds is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of 20 hyaluronic acid and mixtures thereof, the concentration such micelle forming compound being from about 1 to about 5 wt./wt.%. (4)

In another embodiment, suitable for delivery through nasal passages, mixed micellar pharmaceutical 25 formulation is suitably diluted to avoid irritation of the nasal passages.

Another aspect of the present invention provides a mixed micellar pharmaceutical formulation, comprising a pharmaceutical agent in micellar form, water, an alkali 30 metal C8 to C22 alkyl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a

pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one micelle forming compound selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

Yet another aspect of the present invention provides that the mixed micellar aerosol pharmaceutical formulation additionally comprises i) a phenol selected from the group consisting of phenol and methyl phenol in a concentration of from 1 to 10 wt./wt.% of the total formulation, and i) a propellant selected from the group consisting of C1-C2 dialkyl ether, butanes, fluorocarbon propellant, hydrogen-containing fluorocarbon propellant, chlorofluorocarbon propellant, hydrogen-containing chlorofluorocarbon propellant, and mixtures thereof.

In one embodiment, the alkali metal C8 to C22 alkyl sulphate is in a concentration of from 2 to 5 wt./wt.%

of the total formulation.

In another embodiment, the alkali metal C8 to C22 alkyl sulphate is sodium lauryl sulphate.

In another embodiment, the lecithin is saturated or
5 unsaturated, preferably selected from the group consisting of phosphatidylcholine, phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, and lysolecithin.

In yet another embodiment, one of the micelle
10 forming compounds is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, polidocanol alkyl ethers, trihydroxy oxo cholanyl glycine, polyoxyethylene ethers and mixtures thereof, the concentration such absorption
15 enhancing compound being from about 1 to about 5 wt./wt.%.

Preferably, the ratio of pharmaceutical agent, e.g. insulin, to propellant is from 5:95 to 25:75.

In another embodiment, the propellant is selected
20 from the group consisting of tetrafluoroethane, tetrafluoropropane, dimethylfluoropropane, heptafluoropropane, dimethyl ether, n-butane and isobutane.

In yet another embodiment, the mixed micellar
25 pharmaceutical formulation is contained in an aerosol dispenser.

For insulin-containing and some other compositions, the composition may also contains at least one inorganic salt which opens channels in the gastrointestinal tract
30 and may provide additional stimulation to release insulin. Non-limiting examples of inorganic salts are

sodium, potassium, calcium and zinc salts, especially sodium chloride, potassium chloride, calcium chloride, zinc chloride and sodium bicarbonate.

It will be recognized by those skilled in the art
5 that for many pharmaceutical compositions it is usual to add at least one antioxidant to prevent degradation and oxidation of the pharmaceutically active ingredients. It will also be understood by those skilled in the art that colorants, flavouring agents and non-therapeutic
10 amounts of other compounds may be included in the formulation. Typical flavouring agents are menthol and sorbitol.

In one embodiment the antioxidant is selected from the group consisting of tocopherol, deteroxime mesylate,
15 methyl paraben, ethyl paraben and ascorbic acid and mixtures thereof. A preferred antioxidant is tocopherol.

In a preferred embodiment at least one protease inhibitor is added to the formulation to inhibit
20 degradation of the pharmaceutical agent by the action of proteolytic enzymes. Of the known protease inhibitors, most are effective at concentrations of from 1 to 3 wt./wt.% of the formulation.

Non-limiting examples of effective protease
25 inhibitors are bacitracin, soyabean trypsin, aprotinin and bacitracin derivatives, e.g. bacitracin methylene disalicylate. Bacitracin is the most effective of those named when used in concentrations of from 1.5 to 2 wt./wt.%. Soyabean trypsin and aprotinin two may be
30 used in concentrations of about 1 to 2 wt./wt.% of the formulation.

The formulation suitable for delivery through oral mucosal membranes may be in chewable form, in which case it will be necessary to add ingredients suitable for such form. Such ingredients include guar gum, powdered
5 acacia, carrageenin, beeswax and xanthan gum.

The pharmaceutical agent may be selected from a wide variety of macromolecular agents, depending on the disorder being treated, generally with molecular weights greater than about 1000 and especially between about
10 1000 and 2 000 000. Preferred pharmaceutical agents are selected from the group consisting of insulin, heparin, low molecular weight heparin, hirulog, hirugen, huridine, interferons, interleukins, cytokins, mono and polyclonal antibodies, immunoglobins, chemotherapeutic
15 agents, vaccines, glycoproteins, bacterial toxoids, hormones, calcitonins, insulin like growth factors (IGF), glucagon like peptides (GLP-1), large molecule antibiotics, protein based thrombolytic compounds, platelet inhibitors, DNA, RNA, gene therapeutics and
20 antisense oligonucleotides, and small molecule drugs, e.g. opioids, narcotics, analgesics, NSAIDS, steroids, hypnotics, pain killers, morphine and the like.

The present invention also provides a process for making a pharmaceutical composition suitable for
25 delivery through transdermal membranes comprising:
a) preparing a proteinic pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar
30 pharmaceutical agent composition, an alkali metal lauryl sulphate in a concentration of from 1 to 10 wt./wt.% of

the aqueous micellar pharmaceutical agent composition and a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;

- 5 b) slowly adding the micellar proteinic pharmaceutical agent composition to at least one of the absorption enhancing compounds selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,
- 10 octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof, while
- 15 mixing vigorously, to form a mixed micellar composition; wherein the amount of the absorption enhancing compounds are each present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal
- 20 lauryl sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

In one embodiment, the process provides an additional step of adding, while continuing vigorous mixing, at least one absorption enhancing compound

25 different from that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid,

30 linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin,

lysine, polylysine, triolein and mixtures thereof.

In one embodiment the alkali metal lauryl sulphate is sodium lauryl sulphate.

In another embodiment the alkali metal salicylate
5 is sodium salicylate.

In a further embodiment the alkali metal edetate may be selected from the group consisting of disodium edetate and dipotassium edetate.

In yet another embodiment, the formulation has a
10 combinations selected from the group consisting of sodium hyaluronate and unsaturated phospholipid, ii) Phospholipon-H and glycolic acid, and iii) sodium hyaluronate and lecithin.

The present invention also provides a process for
15 making a pharmaceutical composition suitable for delivery by means of an aerosol comprising:

- a) preparing a pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal C8 to C22 alkyl sulphate in a concentration of
20 from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, at least one alkali metal salicylate in a
25 concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;
- b) slowly adding the micellar proteinic pharmaceutical agent composition to at least one of the absorption enhancing compounds selected from the group consisting
30 of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,

- octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine and
- 5 pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, while mixing vigorously, to form a mixed micellar composition; and optionally
- 10 c) an additional step of adding, while continuing vigorous mixing, at least one micelle forming compound different from that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,
- 15 glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linoleic acid, linolenic acid, monoolein, borage oil, evening of primrose oil, glycerin, polyglycerin, lysine, polylysine, triolein, polyoxyethylene ethers and analogues thereof,
- 20 polidocanol alkyl ethers and analogues thereof, and mixtures thereof;
- d) mixing the mixed micellar composition resulting from steps a) to c) with a phenol selected from the group consisting of phenol, m-cresol and mixtures
- 25 thereof; and subsequently
- e) placing the formulation into an aerosol dispenser and charging the dispenser a propellant;
- wherein each of the absorption enhancing compounds are present in a concentration of from 1 to 10 wt./wt.%
- 30 of the total formulation, and the total concentration of alkali metal salicylate, alkali metal C8 to C22 alkyl

sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

The vigorous mixing may be accomplished using high speed stirrers, The vigorous mixing may be accomplished by using high speed stirrers, e.g. magnetic stirrers or propellor stirrers, or by sonication.

In one embodiment, the mixed micellar formulation is formed by sonication of the aqueous micellar pharmaceutical agent composition in the presence of lecithin.

Detailed Description of Preferred Embodiments

The present invention provides an improved method for delivery of macromolecular (high molecular weight) pharmaceutical agents, particularly through the membranes in the nose, mouth, vagina or rectum. The preferred delivery is through oral and nasal cavities. The pharmaceutical agents cover a wide spectrum of agents, including proteins, peptides, hormones, vaccines and drugs. The molecular weights of the macromolecular pharmaceutical agents are preferably above 1000, especially between 1000 and 2 000 000.

For example, hormones which may be administered with the present invention include thyroids, androgens, estrogens, prostaglandins, somatotropins, gonadotropins, erythropoetin, interferons, interleukins, steroids and cytokins. Vaccines which may be administered with the present invention include bacterial and viral vaccines such as vaccines for hepatitis, influenza, tuberculosis, canary pox, chicken pox, measles, mumps, rubella, pneumonia, BCG, HIV and AIDS. Bacterial toxoids which may be administered using the present invention include

diphtheria, tetanus, pseudomonas and mycobactrium tuberculosis. Examples of specific cardiovascular or thromobolytic agents include heparin, hirugen, hirulos and hirudin. Large molecules usefully administered with
5 the present invention include monoclonal antibodies, polyclonal antibodies and immunoglobins.

As will be understood, the concentration of the pharmaceutical agent is an amount sufficient to be effective in treating or preventing a disorder or to
10 regulate a physiological condition in an animal or human. The concentration or amount of pharmaceutical agent administered will depend on the parameters determined for the agent and the method of administration, e.g. oral, nasal. For example, nasal
15 formulations tend to require much lower concentrations of some ingredients in order to avoid irritation or burning of the nasal passages. It is sometimes desirable to dilute an oral formulation up to 10-100 times in order to provide a suitable nasal formulation.

20 The mixed micellar formulation is prepared by first preparing a first micellar composition which contains the pharmaceutically active agents, alkali metal C8 to C22 alkyl sulphate, edetate and alkali metal salicylate. For those compositions intended for administration
25 through the nasal, oral, vaginal or rectal cavities, the first micellar composition is then added to at least one of the absorption enhancing compounds to form a mixed micellar composition. At least one other absorption enhancing compound may also be added subsequently.
30 Preferably the first absorption enhancing compound is lecithin.

When making the aerosol formulation, the phenol and/or m-cresol and/or isotonic agent are then added. The formulation is then put into an aerosol dispenser and the dispenser charged with the propellant.

5 The preferred propellants are hydrogen-containing chlorofluorocarbons, hydrogen-containing fluorocarbons, dimethyl ether and diethyl ether. Even more preferred is hydrofluoroalkane (HFA) 134a (1,1,1,2 tetrafluoroethane).

10 Although the present invention has such wide applicability, the invention is described hereinafter with particular reference to insulin and its analogues, which are used for the treatment of diabetes.

As indicated hereinbefore, the compositions of the
15 present invention require that the pharmaceutical formulation be in mixed micellar form.

In the case of insulin, which is intended for administration through nasal or oral cavities, the first micellar solution may be made by adding a buffer
20 solution to powdered insulin, and then stirring until the powder is dissolved and a clear solution is obtained. A typical buffer solution is an aqueous solution of sodium salicylate and sodium lauryl sulphate and disodium edetate. Typical concentration of sodium
25 salicylate and sodium lauryl sulphate in the aqueous solution are about 3 to 20 wt./wt.% of each compound in the solution. Typically, insulin is present in the micellar solution in an amount which will give a concentration of about 2 to 4 wt./wt.% of the final
30 formulation. Typically the concentration may be about 10 wt./wt.% of the first micellar composition.

The micellar solution is then added slowly to the first absorption enhancing compound, e.g. lecithin while mixing vigorously, e.g. sonicating, to form a mixed micelle liposomal solution. At least one other
5 absorption enhancing compounds selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid,
10 linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein is then added. The mixing may be done with a high speed mixer or sonicator to ensure uniform micelle particle size distribution within
15 the formulation.

Each of the absorption enhancing compounds, when present, is in a concentration of from 1 to 10 wt./wt.% of the total formulation.

Preferred salts of hyaluronic acid are alkali metal
20 hyaluronates, alkaline earth hyaluronates and aluminium hyaluronate. The preferred salt is sodium hyaluronate. The preferred concentration of hyaluronic acid or pharmaceutically acceptable salts of hyaluronic acid is from 1 to 5 wt./wt.% of the total formulation. An even
25 more preferred range is from 1.5 to 3.5 wt./wt.% of the total formulation.

Other ingredients may be added to the mixed micellar solution. For example, flavouring agents, antioxidants, salts, protease inhibitors or other
30 pharmaceutically acceptable compound may be added.

In general the size of the micelle particles in the

solution is about 1 to 10 nm, and preferably from 1 to 5 nm. Such a size distribution ensures effective absorption of the formulation, and therefore the pharmaceutical agent, through the membranes, for example the membranes in the oral and nasal cavities.

The specific concentrations of the essential ingredients can be determined by relatively straightforward experimentation. For absorption through the nasal and oral cavities, it is often desirable to increase, e.g. double or triple, the dosage which is normally required through injection of administration through the gastrointestinal tract.

As will be understood, the amount of each component of the formulation will vary depending on the pharmaceutical agent and the site of application.

Preferred formulations oral or nasal application have the following combinations: i) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H and sodium hyaluronate; ii) sodium lauryl sulphate, sodium salicylate, disodium edetate, lecithin and sodium hyaluronate; iii) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate and evening of primrose oil; iv) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H and bacitracin; v) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H, sodium hyaluronate and bacitracin; and vi) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate, oleic acid and gamma linoleic acid.

For aerosol formulations, the addition of a mixture of phenol and m-cresol is preferred. Such an aerosol

formulation may then be charged to an aerosol dispenser and then charged with a propellant, preferably a non-CFC propellant.

The therapeutic compositions of the present invention may be stored at room temperature or at cold temperature. Storage of proteinic drugs is preferable at a cold temperature to prevent degradation of the drugs and to extend their shelf life.

As indicated hereinbefore, generally, oral and nasal are the favourite sites of the administration but the composition can be applied to the rectal and vaginal mucosa. According to the physiologically active peptide or protein used, the dosage form and the site of administration a specific administration method can be selected.

The composition of this invention is generally prepared as microfine mixed micellar particles (1 to 10 nm or less) by the virtue of its preparation methods used and suitable combinations of absorption enhancer characteristics.

For oral and nasal application, sprays are preferable, but also drops, chewable tablets, chewable gum and other suitable forms may be used. Utilization of atomizer or aerosol spray devices (metered dose inhalers or nebulizers) can be used to further reduce the particle size for effective inhalation from the nasal or oral cavity so the drug may successfully reach to the specific site and be absorbed. It is also possible to utilize a drug delivery system such that an enteric coating is applied to the gelatin capsule to cause the micelles to be released only in the duodenum

or in the proximity of the large intestine and not in the stomach.

The invention is illustrated by reference to the following examples.

5 Example 1

A first experiment was conducted to provide data for comparative purposes. This example does not fall within the scope of the present invention.

6
A solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. To this solution 40 mg (1000 units) of insulin was added and dissolved completely while stirring, to give about 100 units/mL insulin solution.

15 In one set of tests, five healthy non-diabetic human volunteers were tested with insulin, by injection. In another set of tests the volunteers were tested with insulin, taken orally. The volunteers fasted from midnight prior to the test, with no food being taken during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 100 units (1 mL volume per drop, approximately 20 drops) of the above-prepared oral insulin (10 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

25 The average results for the five volunteers, of the first day's trial (sub-cutaneous injection with 10 units) were as follows:

- 24 -

Table I

Time*:	0	15	30	60	75	90	120	150	180
Avg:	5.8	5.8	5.4	5.0	4.6	4.3	3.8	3.6	3.4
Time*:	210	240							
5 Avg:	4.2	4.5							
* time in minutes									

The results for each of the five volunteers, of the second day's trial (oral drops with 100 units) were as follows:

10

Table II

Time*:	0	15	30	60	75	90	120	150	180
Subject Nos:									
1	6.2	5.8	5.2	5.0	4.9	5.0	5.0	4.8	4.7
2	5.8	5.4	5.0	4.7	4.9	4.3	5.0	5.5	5.2
15 3	4.8	4.6	4.3	4.3	4.4	4.6	4.8	4.7	5.2
4	6.6	6.1	5.8	5.5	5.1	4.9	5.0	5.0	5.9
5	6.0	5.8	5.7	5.5	5.1	4.8	4.7	4.9	5.0
Time*:		210	240						
Subject Nos:									
20 1		5.5	6.0						
2		5.8	6.1						
3		5.5	5.1						
4		6.2	6.8						
5		5.9	6.7						
25 *	time in minutes								

These tests indicate that compared to the injection method, oral insulin gives a faster onset of action and lowers blood glucose levels without creating hypoglycaemic condition. Due to the hepatic glucose production, there was a rebound effect. This is believed to be due to the incomplete absorption of insulin.

Example 2

Another experiment, not within the scope of the present invention, was performed for comparative purposes.

- 5 Oral insulin (100 units) was formulated in (Phospholipon-H, 10 mg) without any sodium lauryl sulphate, sodium salicylate, edetate or absorption enhancers, to evaluate its efficacy of blood glucose lowering in a fasted state, for healthy volunteers.
- 10 Volunteers were asked to fast overnight and not have any breakfast prior to dosing. Volunteers were asked to take this oral insulin formulation in their mouth and swallow it. Blood glucose levels were monitored every 15 minutes using Bayer's glucometer
- 15 Elite for 3 hours, and the average results for 5 volunteers are shown in Table III.

Table III

Time*:0	15	30	45	60	75	90	120	150	180
Avg: 5.6	5.8	5.8	5.7	5.7	5.8	5.7	5.7	5.8	5.7

- 20 * time in minutes

This indicates that orally administered insulin with lecithin alone has no effect on blood glucose lowering.

Example 3

- 25 A further experiment, not within the scope of the present invention, was performed for comparative purposes.

- Oral insulin (100 units) was formulated with sodium salicylate and alkali metal edetate (both 5% by wt.) to
- 30 evaluate its efficacy of blood glucose lowering in fasted state in healthy volunteers.

Volunteers were asked to fast overnight and not have any breakfast prior to dosing. Volunteers were asked to take this oral insulin formulation in their mouth and swallow it. Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for 5 volunteers are shown in Table IV.

Table IV

	Time*:0	15	30	45	60	75	90	120	150	180
10	Avg:	5.8	5.8	5.8	5.9	5.8	5.9	5.7	5.9	6.2 6.0

* time in minutes

This indicates that orally administered insulin with sodium salicylate and alkali metal edetate alone has no effect on blood glucose lowering. In addition, this formulation caused irritation and burning sensation, which lasted for several hours.

Example 4

A further experiment, not within the scope of the present invention, was performed for comparative purposes.

Oral insulin (100 units) was formulated using sodium salicylate and alkali metal edetate (both 5% by wt.) with Phospholipon-H (10 mg) and tested on healthy subjects. Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the results are shown in Table V.

Table V

	Time*:	0	15	30	45	60	90	120	180
	Avg:	5.3	5.3	5.3	5.4	5.6	5.7	5.7	5.8

30 * time in minutes

This indicates that orally administered insulin

with sodium salicylate, alkali metal edetate and Phospholipon-H has no effect on blood glucose lowering.

Example 5

Another experiment, not within the scope of the present invention, was performed for comparative purposes.

Oral insulin (50 units) was formulated using only alkali metal lauryl sulphate (5% by wt). Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for four volunteers are shown in Table VI.

Table VI

Time*:	0	15	30	60	90	120	180
Avg:	5.8	5.6	5.4	5.3	5.4	5.4	5.6

* time in minutes

This data shows that orally administered insulin with only alkali metal lauryl sulphate has little metabolic effect on the blood glucose lowering in healthy subjects. This formulation caused substantial burning sensation and irritation in the subjects and lasted for two days.

Example 6

Yet another experiment, within the scope of the present invention, was performed.

Mixed micellar oral insulin (50 units) was formulated using alkali metal lauryl sulphate and sodium salicylate (both 4.4% by wt.) and alkali metal edetate (2.2% by wt) with Phospholipon-H (10 mg) and tested on healthy volunteers.

The method involved mixing the sodium lauryl sulphate, sodium salicylate and alkali metal edetate

with water in a beaker with a magnetic stirrer at medium speed until the ingredients were dissolved, to form buffer solution. Insulin powder was placed in a beaker and to this powder was added the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution obtained. The micellar solution so formed was stored in clean glass bottles and refrigerated.

- 10 Mixed micellar liposomal insulin was then prepared in a glass beaker, in which was placed the Phospholipon-H and a small amount of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10 minutes to ensure complete dissolution of the
- 15 Phospholipon-H. To this solution was added the micellar insulin solution very slowly, drop wise, using glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30 minutes at a high speed to ensure uniform micellar particle size
- 20 distribution.

Samples of the mixed micellar solution were taken orally by the volunteers.

Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for 5 volunteers are shown in Table VII.

Table VII

Time*:	0	15	30	45	60	90	120	150	180
Avg:	6.5	6.1	5.5	5.3	5.3	5.4	5.5	5.5	5.5

30 * time in minutes

This data shows that orally administered insulin

with alkali metal lauryl sulphate combined with the sodium salicylate and alkali metal edetate with Phospholipon-H has a small metabolic effect on blood glucose levels in healthy volunteers.

5 Example 7

An experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

Oral insulin (50 units) was formulated using alkali
10 metal lauryl sulphate and sodium salicylate (both 4.4% by wt.) and alkali metal edetate (2.2% by wt.) with Phospholipon-H (10 mg) and sodium hyaluronate (1.1% by wt). This formulation was tested on healthy subjects under fasting condition.

15 The method involved mixing the sodium lauryl sulphate, sodium salicylate and alkali metal edetate with water in a beaker with a magnetic stirrer at medium speed until the ingredients were dissolved, to form buffer solution. Insulin powder was placed in a beaker
20 and to this powder was added the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution obtained. The micellar solution so formed was stored in clean glass bottles and
25 refrigerated.

Mixed micellar liposomal insulin was then prepared in a glass beaker, in which was placed the Phospholipon-H and a small amount of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10
30 minutes to ensure complete dissolution of the Phospholipon-H. To this solution was added the micellar

insulin solution very slowly, drop wise, using glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30 minutes at a high speed to ensure uniform micellar particle size distribution. The hyaluronate and small amounts of menthol and sorbitol were then added, with continuous stirring.

Samples of the mixed micellar solution were taken orally by the volunteers.

10 Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for 5 volunteers are shown in Table VIII.

Table VIII

15	Time*:	0	15	30	45	60	90	120	150	180
	Avg:	6.5	5.9	5.6	5.4	4.9	5.0	4.9	5.2	5.4

* time in minutes

This data shows that orally administered insulin with alkali metal lauryl sulphate, sodium salicylate, 20 alkali metal edetate, Phospholipon-H and sodium hyaluronate has resulted in lowering of blood glucose levels in healthy subjects better than the above mentioned formulations.

Example 8

25 A further experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g 30 disodium edetate dissolved in 10 mL of water. The solution was added to insulin and mixed, to form

micellar insulin.

Separately, 100 mg of powdered Phosphatidylcholin-H was added to a glass beaker and to this powder was added 10 mL 50% ethanol. The powder was dissolved completely.

5 To this solution 16 mg (400 units) of micellar insulin solution dissolved in 3 mL of the buffer solution to (give 30 units/mL insulin solution) was added slowly with vigorous mixing, to form a mixed micellar solution. To this was added 0.6 mL of sodium hyaluronate and 0.2

10 ml of 2% menthol solution containing 3% sorbitol.

In one set of tests, ten Type II diabetic human volunteers who took insulin, by injection three times a day, were studied. In another set of tests the volunteers were tested with insulin, taken orally. The

15 volunteers fasted from midnight prior to the test, with no food being taken during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the

20 volunteers received 30 units (1 mL volume per drop, approximately 20 drops) of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

25 The results, showing the average for the ten volunteers, were as shown on the following page:

Table IX

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose	Injection
5	(30 units)	(10 units)
0	6.4	6.8
15	5.8	6.9
30	5.4	6.1
45	5.3	5.8
10 60	5.3	5.8
75	5.2	5.8
90	5.2	5.4
105	5.2	5.4
120	5.1	5.2
15 135	5.1	5.1
150	5.2	4.9
165	5.3	4.9
180	5.3	4.8
195	5.4	4.8
20 210	5.4	5.2
225	5.6	5.2
240	5.6	5.4

The results show that the oral insulin formulation of the present invention, at a dosage of three times higher than the injected level, is comparable to the injected insulin.

Example 9

This example illustrates a method for making a mixed micellar formulation according to the present invention.

In a 250 mL capacity glass beaker was added 5 g

sodium lauryl sulphate, 5 g sodium salicylate and 2.5 g edetate. The beaker was placed on the hot plate with a magnetic stirrer. To this dry powder mixture was added 100 mL distilled water and the mixture was stirred, 5 using the magnetic stir bar, at a medium speed until all the powder was dissolved. The buffer solution was stored in a clean glass bottle at room temperature (pH 6.5).

A micellar insulin solution was then prepared in a 10 50 mL capacity glass beaker, into which was placed 11.54 mg insulin powder. To this powder was added 10 mL of the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution 15 obtained. The micellar solution so formed was stored in clean glass bottles and refrigerated.

A 2% menthol solution was then prepared from 100 mg menthol crystals, dissolved in 5 mL ethanol. To this solution was added 5 mg FD & C blue dye. The solution 20 was stirred for 10 minutes and stored in a glass bottle at room temperature.

Mixed micellar liposomal insulin was then prepared in a 50 mL glass beaker, in which was placed 100 mg of phosphatidylcholine (Sigma, type I=EH, hydrogenated). To 25 this powder was added 10 mL of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10 minutes to ensure complete dissolution of the phosphatidylcholine. To this solution was added the micellar insulin solution very slowly, drop wise, using 30 glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30

minutes at a high speed to ensure uniform micellar particle size distribution. To this solution was added 1 mL of the 2% menthol solution and 50 mg sodium hyaluronate. The semi-clear, translucent, light blue colour, liposomal insulin mixed micellar solution (final volume 15 mL) was stored in a clean glass bottle and refrigerated. The solution had a pH of 6.5.

If the phosphatidylcholine powder does not dissolve completely, then heating up to about 45°C may be required, e.g. using a water bath.

It has been found that if the micellar insulin composition is not added slowly, then the mixed micellar formulation will not be formed and the formulation will be gelatinous and sticky.

15 Example 10

The formulation of Example 9 was tested in a manner similar to that indicated in Example 8 except that the formulation of the present invention was administered nasally.

20 On the first day, the ten volunteers each received 10 units insulin injection (regular fast acting, Eli Lilly). On the second day, the volunteers received 20 units of the "oral" insulin of Example 9 (2 times the injection dose). The "oral" insulin was administered as 25 drops (0.4 mL volume per drop, approximately 4 large drops in total, i.e. two drops in each nostril).

The results, showing the average for the ten volunteers, are shown on the following page:

Table X

		Blood glucose levels (mmol/L)	
	Time (minutes)	Nasal Dose	Injection
		(20 units)	(10 units)
5	0	7.4	6.8
	15	6.7	7.0
	30	5.9	6.8
	45	5.3	6.3
	60	5.0	6.3
10	75	5.2	5.8
	90	5.1	5.2
	105	5.0	5.0

Table X (continued)

		Blood glucose levels (mmol/L)	
	Time (minutes)	Nasal Dose	Injection
		(20 units)	(10 units)
15	120	4.6	5.2
	135	4.5	4.2
	150	4.3	4.6
	165	4.3	4.0
	180	4.8	4.1
20	195	5.3	4.3
	210	5.4	4.5
	225	5.7	4.7
	240	5.6	5.0

The results show that the nasal insulin formulation of the present invention, at a dosage of twice the injected level, is comparable to the injected insulin.

Example 11

30 The formula of Example 9 was taken and tests performed to determine the insulin action on meal

glucose on healthy volunteers.

Usually, diabetic patients take an insulin injection 30 minutes prior to a meal, because injected insulin takes a long time to take effect. Injected
5 insulin is slowly absorbed into bloodstream within 60 minutes and has metabolic effect on meal glucose levels.

The mixed micellar formulation of Example 9 was test 1 in healthy volunteers under controlled conditions to determine the oral insulin effect on meal glucose
10 when compared to injected insulin.

In one set of tests, ten healthy non-diabetic human volunteers were tested with insulin, by injection. In another set of tests the volunteers were tested with insulin, taken orally. The volunteers fasted from
15 midnight prior to the tests, with food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units
20 insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 30 units of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by
25 Bayer's Glucometer Elite. The results are shown on the following page:

Table XI

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose	Injection
	(30 units)	(10 units)
5		
0	5.7	5.5
15	5.2	5.6
30	5.0	5.4
45	5.3	5.4
10 60	5.4	5.6
75	6.3	6.6
90	6.9	7.0
105	6.0	5.9
120	5.8	5.6
15 135	5.5	5.1
150	5.1	4.8
165	4.9	4.6
180	4.8	4.3

The results indicate that the oral insulin helps
 20 control meal glucose levels in healthy volunteers when
 compared to injected insulin.

Example 12

The mixed micellar formulation of Example 9 was
 tested in diabetic volunteers under controlled
 25 conditions to determine the oral insulin effect on meal
 glucose when compared to injected insulin.

In one set of tests, ten Type II diabetic human
 volunteers who took insulin, by injection three times a
 day, were studied. In another set of tests the
 30 volunteers were tested with insulin, taken orally. The
 volunteers fasted from midnight prior to the tests, with

food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units
5 insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 30 units of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by
10 Bayer's Glucometer Elite.

The average results for the 10 volunteers were as follows:

Table XII

		Blood glucose levels (mmol/L)	
15	Time (minutes)	Oral Dose	Injection
		(30 units)	(10 units)
	0	8.8	8.7
	15	8.1	8.8
	30	8.0	8.9
20	45	8.4	10.1
	60	10.2	11.8
	75	11.8	11.8
	90	12.3	12.2
	105	10.8	11.2
25	120	9.6	10.4
	135	8.1	8.4
	150	6.9	7.3
	165	6.2	6.5
	180	4.8	4.3

30 The results indicate that oral insulin helps to control meal glucose levels in diabetic patients when

compared to injected insulin.

Example 13

A chewable gum insulin formulation was prepared by vigorously stirring the liposomal insulin mixed micellar solution of Example 9 while adding guar gum, beeswax, powdered acacia, oleic acid, gamma-linoleic acid and sorbitol. For each 30 units of insulin, the mixture contained 100 mg guar gum, 50 mg beeswax, 50 mg powdered acacia, 100 mg oleic acid, 100 mg gamma-linoleic acid and 1 mL 3% sorbitol in ethanol solution. The mixture was then poured into a flat tray coated with polytetrafluoroethylene until the mixture was about 10 mm deep. The mixture then solidified and after solidification was cut into sticks about 1 cm by 3 cm. Each stick contained about 30 units insulin.

The mixed micellar formulation in chewable stick form was tested in diabetic volunteers under controlled conditions to determine the oral insulin effect on meal glucose when compared to injected insulin.

In one set of tests, five Type II diabetic human volunteers who took insulin, by injection three times a day, were studied. In another set of tests the volunteers were tested with the chewable gum insulin, taken orally. The volunteers fasted from midnight prior to the tests, with food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the

volunteers received 30 units of the above-prepared chewable gum oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

- 5 The average results for the five volunteers were as follows:

Table XIII

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose	Injection
	(30 units)	(10 units)
10 0	9.1	8.8
15	9.3	8.2
30	9.3	8.0
45	10.2	8.4
15 60	11.2	9.2
75	12.1	10.3
90	12.9	11.8
105	13.2	11.6
120	12.8	11.0
20 135	12.2	10.2
150	11.6	9.6
165	11.0	9.5
180	10.6	9.1
195	10.0	8.7
25 210	9.5	8.2
225	8.8	8.0
240	8.2	7.5

Example 14

Another experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. The solution was added to 8 mg (200 units) insulin and 5 mixed, to form micellar insulin.

To this micellar solution were added 0.2 g bacitracin and 0.5 g evening of primrose oil and the solution was mixed vigorously to form a mixed micellar insulin solution (about 20 units/mL).

10 Six human volunteers were studied. The volunteers fasted from midnight prior to the test, with no food being taken during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, 15 available from Eli Lilly). On the second day, the volunteers received 20 units of the above-prepared oral insulin (twice the injection dose). In both tests, blood glucose levels were monitored at intervals by Bayer's Glucometer Elite.

20 The results, showing the average for the six volunteers, were as follows:

Table XIV

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose	Injection
25	(20 units)	(10 units)
0	8.8	7.9
15	8.4	7.9
30	8.1	8.2
45	7.4	8.3
30 60	6.3	7.6
90	5.1	6.2

120	5.0	5.2
150	4.8	4.6
180	5.1	3.9
210	5.3	4.4
5 240	5.6	5.2

The results show that the oral insulin formulation of the present invention, at a dosage of twice the injected level, is comparable to the injected insulin.

Example 15

10 A further experiment was performed to show another method of making the mixed micellar formulation of the present invention.

In a 250 mL round bottom flask was added 100 mg of saturated lecithin powder (Phospholipon-90H) purchased
15 from the American Lecithin Co. To this powder was added 5 mL of absolute ethanol (USP grade). The flask was then attached to a rotary evaporator equipped with the vacuum pump and nitrogen inlet for inert atmosphere condition to minimize oxidation of the lecithin. The
20 flask was rotated at 100-150 rpm under vacuum. The solution in the flask was heated to 60°C by means of water bath to dissolve the powder completely. After complete dissolution of the powder, heating was stopped and the rotation speed was increased to 300 rpm, under
25 vacuum in nitrogen atmosphere until the alcohol evaporated completely, leaving a uniform film on the side of the flask. The rotation was continued for at least 30 minutes to ensure uniform coating of film on the wall and complete solvent removal. After 30
30 minutes the rotation was stopped and the vacuum was released.

To this flask was added micellar insulin solution which had been prepared from an aqueous solution of insulin, sodium lauryl sulphate, sodium salicylate and disodium edetate. The flask was shaken with the help of shaker plate. Shaking was continued for at least 30 minutes and then the solution was sonicated with a high frequency sonicating probe for another 60 minutes in order to form small uniform mixed micelles. The mixed micelles so obtained were analyzed by Malvern Zeta (trade mark) particle size distribution measurement equipment equipped with the laser light scattering device. The mixed micelles particle size distribution obtained by this method was between 2 and 9nm. To this solution was added 1 mL of 2% menthol solution and 50 mg sodium hyaluronate. The semi-clear, translucent, light blue colour solution (final volume 10 mL) was stored in a clean glass bottle and refrigerated. The solution had a pH of 6.5.

Example 16

Another experiment, within the scope of the present invention, was performed.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. The solution was added to 8 mg (200 units) insulin and mixed, to form micellar insulin.

To this micellar solution were added 0.5 g borage oil and the solution was mixed vigorously to form a mixed micellar insulin solution (about 20 units/mL).

CLAIMS:

1. A process for making a pharmaceutical composition suitable for delivery through mucosal membranes comprising:
 - 5 a) preparing a pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, an alkali metal C8 to C22 alkyl sulphate in
10 a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition and a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;
 - 15 b) slowly adding the micellar proteinic pharmaceutical agent composition, while mixing, to at least one absorption enhancing compound, while continuing to mix vigorously, said absorption enhancing compounds being selected from the group consisting of lecithin,
20 hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine
25 and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to
30 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal

C8 to C22 alkyl sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

2. A process according to Claim 1 wherein there is an
5 additional step of adding, while continuing mixing, at least one absorption enhancing compound different to that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,
10 octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof.
- 15 3. A process according to Claim 1 wherein the absorption enhancing compound in step b) is selected from the group consisting of saturated phospholipid, unsaturated phospholipid, phosphatidylcholine, phosphatidyl serine, sphingomyelin,
20 phosphatidylethanolamine, cephalin, lecithin, lysolecithin and mixtures thereof.
4. A process according to Claim 1 wherein one of the absorption enhancing compounds is lecithin and another absorption enhancing compound is selected from the group
25 consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid and mixtures thereof, the concentration such absorption enhancing compound being from about 1 to about 5 wt./wt.%.
5. A process according to Claim 1 wherein the micellar
30 absorption enhancing compounds comprise combinations selected from the group consisting of i) saturated

phospholipid and sodium hyaluronate, ii) saturated phospholipid and glycolic acid, iii) lecithin and sodium hyaluronate and iv) saturated phospholipid, glycolic acid and lactic acid.

- 5 6. A process according to Claim 1 wherein the proteinic pharmaceutical agent is selected from the group consisting of insulin, heparin, so-called low molecular weight heparin, hirulog, hirugen, hirudin, interferons, interleukins, cytokines, mono and
- 10 polyclonal antibodies, chemotherapeutic agents, vaccines, glycoproteins, bacterial toxoids, hormones, calcitonins, insulin like growth factors (IGF), glucagon like peptides (GLP-1), large molecule antibiotics, protein based thrombolytic compounds, platelet
- 15 inhibitors, DNA, RNA, gene therapeutics, antisense oligonucleotides, opioids, narcotics, analgesics, NSAIDS, steroids, hypnotics, pain killers and morphine.

7. A process according to Claim 1 wherein in step b) the micellar proteinic pharmaceutical agent composition
- 20 is added to lecithin, with sonication, to form a mixed micellar composition; and

- c) while continuing to mix, adding at least one absorption enhancing compound selected from the group consisting of hyaluronic acid, pharmaceutically
- 25 acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin,
- 30 lysine, polylysine, triolein and mixtures thereof;

wherein the amount of lecithin and the absorption

enhancing compound are each present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal C8 to C22 alkyl sulphate, edetate and
5 absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

8. A process according to Claim 1 wherein the absorption enhancing compound is formed into a film prior to the addition of the micellar pharmaceutical
10 agent composition.

9. A process according Claim 1 wherein subsequent to the addition of the micellar pharmaceutical agent composition a second absorption enhancing compound is added, said second absorption enhancing compound being
15 different from the absorption enhancing compound previously used.

10. A process according to Claim 1 a phenol selected from the group consisting of phenol, methyl phenol and mixtures thereof are added to the micellar formulation
20 and the resulting formulation placed in a container, and the container is subsequently charged with a propellant.

11. A process according to Claim 10 wherein the propellant is selected from the group consisting of tetrafluoroethane, tetrafluoropropane,
25 dimethylfluoropropane, heptafluoropropane, dimethyl ether, n-butane and isobutane.

12. A process according to Claim 1 wherein the pharmaceutical agent is insulin.

13. A process according to Claim 11 wherein the
30 pharmaceutical agent is insulin.

14. A mixed micellar pharmaceutical formulation

comprising a pharmaceutical agent in micellar form, water, an alkali metal C8 to C22 alkyl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a pharmaceutically acceptable edetate in a
5 concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one absorption forming compound, said absorption forming compounds being
10 selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of
15 primrose oil, menthol, trihydroxy oxo cholanylglycine and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing
20 compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

15. A mixed micellar pharmaceutical formulation
25 according to Claim 14, in which one of the absorption enhancing compounds is lecithin.

16. A formulation according to Claim 14 wherein the alkali metal C8 to C22 alkyl sulphate is sodium lauryl sulphate and the alkali metal salicylate is sodium
30 salicylate.

17. A formulation according to Claim 15 wherein the

lecithin is selected from the group consisting of saturated phospholipid, unsaturated phospholipid, phosphatidylcholine, phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, lysolecithin and

5 mixtures thereof.

18. A formulation according to Claim 15 wherein the other absorption enhancing compound is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid and
10 mixtures thereof, the concentration such absorption enhancing compound being from about 1 to about 5 wt./wt.%.
19. A formulation according to Claim 14 wherein the

formulation comprises combinations selected from the
15 group consisting of i) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated phospholipid and sodium hyaluronate; ii) sodium lauryl sulphate, sodium salicylate, disodium edetate, lecithin and sodium hyaluronate; iii) sodium lauryl sulphate, sodium
20 salicylate, disodium edetate, sodium hyaluronate and evening of primrose oil; iv) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated phospholipid and bacitracin; v) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated
25 phospholipid, sodium hyaluronate and bacitracin; and vi) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate, oleic acid and gamma linoleic acid.

20. A formulation according to Claim 14 wherein the
30 pharmaceutical agent is selected from the group consisting of insulin, heparin, so-called low molecular

weight heparin, hirulog, hirugen, huridine, interferons,
interleukins, cytokins, mono and polyclonal antibodies,
chemotherapeutic agents, vaccines, glycoproteins,
bacterial toxoids, hormones, calcitonins, insulin like
5 growth factors (IGF), glucagon like peptides (GLP-1),
large molecule antibiotics, protein based thrombolytic
compounds, platelet inhibitors, DNA, RNA, gene
therapeutics antisense oligonucleotides, opioids,
narcotics, analgesics, NSAIDS, steroids, hypnotics, pain
10 killers and morphine.

21. A formulation according to Claim 14 wherein the
pharmaceutical agent is insulin.

22. A formulation according to Claim 21 in which the
absorption enhancing compounds are lecithin and a second
15 absorption enhancing compound selected from the group
consisting of hyaluronic acid, pharmaceutically
acceptable salts of hyaluronic acid and mixtures
thereof.

23. A formulation according to Claim 14 wherein the
20 formulation additionally comprises a phenol selected
from the group consisting of phenol, methyl phenol and
mixtures thereof.

24. A formulation according to Claim 23 wherein the
formulation is contained in an aerosol container and the
25 container is charged with a propellant.

25. A formulation according to Claim 24 wherein the
propellant is selected from the group consisting of
tetrafluoroethane, tetrafluoropropane,
dimethylfluoropropane, heptafluoropropane, dimethyl
30 ether, n-butane and isobutane.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 99/00106

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K38/28 A61K9/107

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of documents with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 36352 A (MODI PANKAJ ;CHANDARANA SUBASH (CA)) 21 November 1996 see page 3, line 2-27 see page 5, line 7-15 see page 5, line 35 - page 6, line 25 see page 7, line 4-5 see claims	1-7,9, 12-22
A	WO 96 19197 A (ASTRA AB ;BAECKSTROEM KJELL (SE); DAHLBAECK MAGNUS (SE); JOHANSSON) 27 June 1996 see page 2, line 15-24 see page 3, line 26-30 see page 4, line 10-27 see page 5, line 22 - page 6, line 14 see examples see claims 1,8-12,15-17,24-26,43	1,6, 10-14, 24,25
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 June 1999

Date of mailing of the international search report

10/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

La Gaetana, R

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/CA 99/00106

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>YAMAMOTO A ET AL: "Pulmonary absorption enhancement of peptides by absorption enhancers and protease inhibitors" JOURNAL OF CONTROLLED RELEASE, vol. 41, no. 1, 1 August 1996, page 57-67 XP004037571 see abstract see pages 60-62, "Results", paragraphs 3.1-3.3 see page 64, column 2, line 24-page 65, column 1, line 5</p>	1, 14
A	<p>WO 94 26302 A (GENENTECH INC) 24 November 1994 see page 3, line 2-22 see page 6, line 18-27 see page 7, line 4-23 see claims 1, 11, 12, 14</p>	1, 10, 11, 14, 23, 24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 99/00106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9636352 A	21-11-1996	US 5653987 A	05-08-1997
		AU 5642396 A	29-11-1996
		CA 2210996 A	21-11-1996
		EP 0813421 A	29-12-1997
WO 9619197 A	27-06-1996	AU 702879 B	11-03-1999
		AU 4359196 A	10-07-1996
		BR 9510501 A	13-01-1998
		CA 2206736 A	27-06-1996
		CN 1171046 A	21-01-1998
		CZ 9701945 A	15-10-1997
		EP 0797431 A	01-10-1997
		FI 972657 A	19-06-1997
		HU 77701 A	28-07-1998
		JP 10510827 T	20-10-1998
		NO 972781 A	16-06-1997
		PL 320824 A	10-11-1997
		SK 81397 A	05-11-1997
		ZA 9510752 A	24-06-1996
WO 9426302 A	24-11-1994	AT 173632 T	15-12-1998
		AU 683469 B	13-11-1997
		AU 6782294 A	12-12-1994
		CA 2159602 A	24-11-1994
		CN 1122575 A	15-05-1996
		CZ 9502955 A	17-04-1996
		DE 69414841 D	07-01-1999
		DE 69414841 T	02-06-1999
		EP 0697887 A	28-02-1996
		ES 2126114 T	16-03-1999
		FI 955136 A	27-10-1995
		HU 72980 A	28-06-1996
		JP 8510242 T	29-10-1996
		NO 954544 A	10-11-1995
		NZ 266354 A	24-06-1997
		PL 311639 A	04-03-1996
		SK 137895 A	05-02-1997
		ZA 9402955 A	30-10-1995

CORRECTED
VERSION*

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 38/28, 9/107	A1	(11) International Publication Number: WO 99/40932 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/CA99/00106 (22) International Filing Date: 5 February 1999 (05.02.99) (30) Priority Data: 09/021,114 10 February 1998 (10.02.98) US 09/216,733 21 December 1998 (21.12.98) US (71)(72) Applicant and Inventor: MODI, Pankaj (CA/CA); 519 Golf Links Road, Ancaster, Ontario L9G 4X6 (CA). (74) Agent: BARRIGAR & MOSS; Suite 901, 2 Robert Speck Parkway, Mississauga, Ontario L4Z 1H8 (CA).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: MIXED MICELLAR PHARMACEUTICAL DELIVERY SYSTEM AND METHOD OF PREPARATION (57) Abstract A mixed micellar pharmaceutical formulation includes a micellar proteinic pharmaceutical agent, an alkali metal C8 to C22 alkyl sulphate, alkali metal salicylate, a pharmaceutically acceptable edetate and at least one absorption enhancing compounds. The absorption enhancing compounds are selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening primrose oil, trihydroxy oxo cholanyl glycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof. The amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.		

* (Referred to in PCT Gazette No. 13/2000, Section II)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MIXED MICELLAR PHARMACEUTICAL DELIVERY SYSTEM AND METHOD OF PREPARATION

Field of the Invention

The present invention relates to an improved
5 delivery system for the administration of large-molecule
pharmaceuticals, e.g. peptidic drugs, vaccines and
hormones. In particular it relates to pharmaceuticals
which may be administered through the oral and nasal
membranes.

10 Background to the Invention

In spite of significant efforts in academic and
commercial laboratories, major breakthroughs in oral
peptide and protein formulation have not been achieved.
Relatively little progress has been made in reaching the
15 target of safe and effective oral formulations for
peptides and proteins. The major barriers to developing
oral formulations for proteins and peptides include poor
intrinsic permeability, lumenal and cellular enzymatic
degradation, rapid clearance, and chemical stability in
20 the gastrointestinal (GI) tract. Pharmaceutical
approaches to address these barriers, which have been
successful with traditional small, organic drug
molecules, have not readily translated into effective
peptide and protein formulations. Although the
25 challenges are significant, the potential therapeutic
benefits remain high especially in the field of diabetes
treatment using insulin.

Scientists have explored various administration
routes other than the injection for proteins and
30 peptides. These routes include oral, intranasal, rectal,
vaginal cavities for the effective delivery of large
molecules. Out of the above four mentioned routes oral

and nasal cavities have been of greatest interest to scientists. Both the oral and nasal membranes offer advantages over other routes of administration. For example, drugs administered through these membranes have

5 a rapid onset of action, provide therapeutic plasma levels, avoid first pass effect of hepatic metabolism, and avoid exposure of the drug to hostile GI environment. Additional advantages include easy access to the membrane sites so that the drug can be applied,

10 localized and removed easily. Further, there is a good potential for prolonged delivery of large molecules through these membranes.

The oral routes have received far more attention than has the other routes. The sublingual mucosa

15 includes the membrane of ventral surface of the tongue and the floor of the mouth whereas the buccal mucosa constitutes the lining of the cheek. The sublingual mucosa is relatively permeable thus giving rapid absorption and acceptable bioavailability of many drugs.

20 Further, the sublingual mucosa is convenient, acceptable and easily accessible. This route has been investigated clinically for the delivery of a substantial number of drugs.

The ability of molecules to permeate through the

25 oral mucosa appears to be related to molecular size, lipid solubility and peptide protein ionization. Small molecules, less than 1000 daltons appear to cross mucosa rapidly. As molecular size increases, the permeability decreases rapidly. Lipid soluble compounds are more

30 permeable than non-lipid soluble molecules. Maximum absorption occurs when molecules are un-ionized or

neutral in electrical charges. Therefore charged molecules present the biggest challenges to absorption through the oral mucosae.

Most proteinic drug molecules are extremely large molecules with molecular weight exceeding 6000 daltons. These large molecules have very poor lipid solubility and are practically impermeable. Substances that facilitate the absorption or transport of large molecules (>2000 daltons) across biological membranes are known as the enhancers, (Lee et al., Critical Reviews in Therapeutic drug Carrier Systems, 8, 91, 1991; Lee et al., Critical Reviews in Therapeutic drug Carrier Systems, 8, 115, 1991, 1992). Enhancers may be characterized as chelators, bile salts, fatty acids, synthetic hydrophilic and hydrophobic compounds, and biodegradable polymeric compounds.

Various mechanisms of action of enhancers have been proposed. These mechanisms of action, at least for protein and peptidic drugs include (1) reducing viscosity and/or elasticity of mucous layer, (2) facilitating transcellular transport by increasing the fluidity of the lipid bilayer of membranes, and (3) increasing the thermodynamic activity of drugs (Critical Rev, 117-125, 1991, 1992).

Many enhancers have been tested so far and some have found to be effective in facilitating mucosal administration of large molecule drugs. However, hardly any penetration enhancing products have reached the market place. Reasons for this include lack of a satisfactory safety profile respecting irritation, lowering of the barrier function, and impairment of the

mucocilliary clearance protective mechanism. The main factor to be considered in the use of enhancers especially related to bile salts, and some protein solubilizing agents is extremely bitter and unpleasant taste. This makes their use almost impossible for human consumption on a daily basis. Several approaches were utilized to improve the taste of the bile salts based delivery systems, but none one of them are commercially acceptable for human consumption to date. Among the approaches utilized includes patch for buccal mucosa, bilayer tablets, controlled release tablets, use of protease inhibitors, buccally administered film patch devices, and various polymer matrices.

The basic problem associated with the above technologies is the use of large quantities of bile acids and their salts to promote the transport of the large molecules through membranes in the form of localized delivery system using patches or tablets. In spite of using protease inhibitors and polymer coatings the technologies failed to deliver proteinic drugs in the required therapeutic concentrations. Further, the problem is compounded because of the localized site effect of the patch which resulted in severe tissue damage in the mouth. Most attempts were made to deliver large molecules via the oral, nasal, rectal, and vaginal routes using single bile acids or enhancing agents in combination with protease inhibitors and biodegradable polymeric materials. However, it is extremely difficult to achieve therapeutic levels of proteinic drugs using these formulations. As single enhancing agents fails to loosen tight cellular junctions in the oral, nasal,

rectal and vaginal cavities for a required period of time to allow passage of large molecules through the mucosal membranes without further degradation. This problem makes it impractical to use the above mentioned systems for a commercial purpose.

In order to overcome the above mentioned problem of the bitter taste, irritation and the penetration of large molecules through the sublingual, buccal and GI tract mucosal lining, a system has now been designed where protein drug was encapsulated in mixed micelles made up of combination of enhancers, e.g. yolk proteins (lecithins). This system allows opening of the paracellular junctions (tight junctions) in oral as well as in GI tract by GI motility movement with high degree of protease activity preserved and protecting molecules from premature degradation in the hostile acidic and proteolytic GI environment.

It is believed that the mixed micelles encapsulate molecules with high degree of efficiency (>90% encapsulation). These mixed micelles are extremely small in the size (1 nm to 10 nm), and are smaller than the pores of the membranes in the oral cavity or the GI tract. It is therefore believed that the extremely small size of mixed micelles helps encapsulated molecules penetrate efficiently through the mucosal membranes of the oral cavity.

The absorption of proteins and peptides is believed to be enhanced by the diffusion of large molecules entrapped in the mixed micellar form through the aqueous pores and the cell structure perturbation of the tight paracellular junctions.

The amount of physiologically peptide or protein in the compositions of this invention is typically a quantity that provides an effective amount of the drug to produce the physiological activity (therapeutic plasma level) for which peptide or protein is being administered. In consideration of the fact that the bioavailability of any active substance can never be 100%, that is to say the administered dose of the active drug is not completely absorbed, it is preferable to incorporate slightly larger amount than the desired dosage. Where the dosage form is a spray (aerosol) or the like which is repeatedly dispensed from the same container, it is recommendably so arranged that the unit dose will be slightly greater than the desired dose. It should be understood that dosage should vary with species of warm blood animals such as man, domestic animals, and their body weights. Although the composition of this invention is prepared as the microfine droplets (1 to 10 nm or less) by the virtue of its preparation methods used and suitable combinations of enhancer compound characteristics. The utilization of atomizer or aerosol spray devices (metered dose inhalers or nebulizers) may be useful to further a sufficient reduction of particle size for effective inhalation from the nasal or oral cavity so the drug may successfully absorbed or reach to the specific site.

The therapeutic composition of the present invention can be stored at room temperature or at cold temperature. Storage of proteinic drugs is preferable at the cold temperature to prevent the degradation of the drugs and to extend their shelf life. While the mixed

micellar therapeutic composition of the invention is applied to the mucosal membranes, the sites of administration may be the same as those used for the usual mucosal therapeutic preparation. Generally, oral, 5 transdermal and nasal are the favourite sites of the administration but the composition can be applied to the rectal and vaginal mucosa. According to the physiologically active peptide or protein used, the dosage form and the site of administration, a specific 10 administration method can be selected.

As used herein, the term "edetate" is used herein to refer to pharmaceutically acceptable salts of ethylenediaminetetraacetic acid.

It has also been found that improvements in 15 penetration and absorption of mixed micellar formulations can be achieved by mixing the mixed micellar formulation with propellants such as tetrafluoroethane, heptafluoroethane, dimethylfluoropropane, tetrafluoropropane, butane, 20 isobutane, dimethyl ether and other non-CFC and CFC propellants. Preferably they are delivered through metered dose spray devices. Metered dose inhalers are known and are a popular pulmonary drug delivery form for some drugs. The present formulation, including the 25 propellant is intended to improve the quality of absorption, stability and performance of many formulations. The compositions have been selected to give enhancement in the penetration through pores, and facilitate absorption of the drugs to reach therapeutic 30 levels in the plasma. The present formulation may be absorbed buccally, by ensuring that the person does not

inhale the formulation as it is sprayed. One of the other benefits of using an atomizer or inhaler is that the potential for contamination is minimized because the devices are self contained.

5 Summary of the Invention

Accordingly the present invention provides a mixed micellar pharmaceutical formulation, having a pH of between 6.0 and 7.0 comprising a proteinic pharmaceutical agent in micellar form, water, an alkali metal lauryl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one micelle forming compound selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

In an embodiment, the alkali metal lauryl sulphate, the edetate and the alkali metal salicylate are each in a concentration of from 2 to 5 wt./wt.% of the total

formulation.

In one embodiment, the edetate is an alkali metal edetate. Preferably the alkali metal edetate is be selected from the group consisting of disodium edetate,
5 dipotassium edetate, and combinations thereof.

In another embodiment, the alkali metal lauryl sulphate is sodium lauryl sulphate.

In a further embodiment, the alkali metal salicylate is sodium salicylate.

10 In another embodiment, the lecithin is selected from the group consisting of saturated phospholipid, e.g. Phospholipon-H (trade mark) saturated phospholipid, unsaturated phospholipid, e.g. Phospholipon-G (trade mark) unsaturated phospholipid, phosphatidylcholine,
15 phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, and lysolecithin.

In one embodiment, one of the absorption enhancing compounds is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of
20 hyaluronic acid and mixtures thereof, the concentration such micelle forming compound being from about 1 to about 5 wt./wt.%.
6

In another embodiment, suitable for delivery through nasal passages, mixed micellar pharmaceutical
25 formulation is suitably diluted to avoid irritation of the nasal passages.

Another aspect of the present invention provides a mixed micellar pharmaceutical formulation, comprising a pharmaceutical agent in micellar form, water, an alkali
30 metal C8 to C22 alkyl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a

- pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one micelle forming compound selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.
- Yet another aspect of the present invention provides that the mixed micellar aerosol pharmaceutical formulation additionally comprises i) a phenol selected from the group consisting of phenol and methyl phenol in a concentration of from 1 to 10 wt./wt.% of the total formulation, and ii) a propellant selected from the group consisting of C1-C2 dialkyl ether, butanes, fluorocarbon propellant, hydrogen-containing fluorocarbon propellant, chlorofluorocarbon propellant, hydrogen-containing chlorofluorocarbon propellant, and mixtures thereof.
- In one embodiment, the alkali metal C8 to C22 alkyl sulphate is in a concentration of from 2 to 5 wt./wt.%

of the total formulation.

In another embodiment, the alkali metal C8 to C22 alkyl sulphate is sodium lauryl sulphate.

In another embodiment, the lecithin is saturated or
5 unsaturated, preferably selected from the group consisting of phosphatidylcholine, phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, and lysolecithin.

In yet another embodiment, one of the micelle
10 forming compounds is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, polidocanol alkyl ethers, trihydroxy oxo cholanyl glycine, polyoxyethylene ethers and mixtures thereof, the concentration such absorption
15 enhancing compound being from about 1 to about 5 wt./wt.%.
5 wt./wt.%.

Preferably, the ratio of pharmaceutical agent, e.g. insulin, to propellant is from 5:95 to 25:75.

In another embodiment, the propellant is selected
20 from the group consisting of tetrafluoroethane, tetrafluoropropane, dimethylfluoropropane, heptafluoropropane, dimethyl ether, n-butane and isobutane.

In yet another embodiment, the mixed micellar
25 pharmaceutical formulation is contained in an aerosol dispenser.

For insulin-containing and some other compositions, the composition may also contains at least one inorganic salt which opens channels in the gastrointestinal tract
30 and may provide additional stimulation to release insulin. Non-limiting examples of inorganic salts are

sodium, potassium, calcium and zinc salts, especially sodium chloride, potassium chloride, calcium chloride, zinc chloride and sodium bicarbonate.

It will be recognized by those skilled in the art
5 that for many pharmaceutical compositions it is usual to add at least one antioxidant to prevent degradation and oxidation of the pharmaceutically active ingredients. It will also be understood by those skilled in the art that colorants, flavouring agents and non-therapeutic
10 amounts of other compounds may be included in the formulation. Typical flavouring agents are menthol and sorbitol.

In one embodiment the antioxidant is selected from the group consisting of tocopherol, deteroxime mesylate,
15 methyl paraben, ethyl paraben and ascorbic acid and mixtures thereof. A preferred antioxidant is tocopherol.

In a preferred embodiment at least one protease inhibitor is added to the formulation to inhibit
20 degradation of the pharmaceutical agent by the action of proteolytic enzymes. Of the known protease inhibitors, most are effective at concentrations of from 1 to 3 wt./wt.% of the formulation.

Non-limiting examples of effective protease
25 inhibitors are bacitracin, soyabean trypsin, aprotinin and bacitracin derivatives, e.g. bacitracin methylene disalicylate. Bacitracin is the most effective of those named when used in concentrations of from 1.5 to 2 wt./wt.%. Soyabean trypsin and aprotinin two may be
30 used in concentrations of about 1 to 2 wt./wt.% of the formulation.

The formulation suitable for delivery through oral mucosal membranes may be in chewable form, in which case it will be necessary to add ingredients suitable for such form. Such ingredients include guar gum, powdered
5 acacia, carrageenin, beeswax and xanthan gum.

The pharmaceutical agent may be selected from a wide variety of macromolecular agents, depending on the disorder being treated, generally with molecular weights greater than about 1000 and especially between about
10 1000 and 2 000 000. Preferred pharmaceutical agents are selected from the group consisting of insulin, heparin, low molecular weight heparin, hirulog, hirugen, hiridine, interferons, interleukins, cytokins, mono and polyclonal antibodies, immunoglobins, chemotherapeutic
15 agents, vaccines, glycoproteins, bacterial toxoids, hormones, calcitonins, insulin like growth factors (IGF), glucagon like peptides (GLP-1), large molecule antibiotics, protein based thrombolytic compounds, platelet inhibitors, DNA, RNA, gene therapeutics and
20 antisense oligonucleotides, and small molecule drugs, e.g. opioids, narcotics, analgesics, NSAIDS, steroids, hypnotics, pain killers, morphine and the like.

The present invention also provides a process for making a pharmaceutical composition suitable for
25 delivery through transdermal membranes comprising:
a) preparing a proteinic pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar
30 pharmaceutical agent composition, an alkali metal lauryl sulphate in a concentration of from 1 to 10 wt./wt.% of

the aqueous micellar pharmaceutical agent composition and a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;

- 5 b) slowly adding the micellar proteinic pharmaceutical agent composition to at least one of the absorption enhancing compounds selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,
- 10 octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof, while
- 15 mixing vigorously, to form a mixed micellar composition; wherein the amount of the absorption enhancing compounds are each present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal
- 20 lauryl sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

In one embodiment, the process provides an additional step of adding, while continuing vigorous mixing, at least one absorption enhancing compound

25 different from that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid,

30 linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin,

lysine, polylysine, triolein and mixtures thereof.

In one embodiment the alkali metal lauryl sulphate is sodium lauryl sulphate.

In another embodiment the alkali metal salicylate
5 is sodium salicylate.

In a further embodiment the alkali metal edetate may be selected from the group consisting of disodium edetate and dipotassium edetate.

In yet another embodiment, the formulation has a
10 combinations selected from the group consisting of sodium hyaluronate and unsaturated phospholipid, ii) Phospholipon-H and glycolic acid, and iii) sodium hyaluronate and lecithin.

The present invention also provides a process for
15 making a pharmaceutical composition suitable for delivery by means of an aerosol comprising:

- a) preparing a pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal C8 to C22 alkyl sulphate in a concentration of
20 from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, at least one alkali metal salicylate in a
25 concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;
- b) slowly adding the micellar proteinic pharmaceutical agent composition to at least one of the absorption enhancing compounds selected from the group consisting
30 of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,

- octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine and
- 5 pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, while mixing vigorously, to form a mixed micellar composition; and optionally
- 10 c) an additional step of adding, while continuing vigorous mixing, at least one micelle forming compound different from that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid,
- 15 glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linoleic acid, linolenic acid, monoolein, borage oil, evening of primrose oil, glycerin, polyglycerin, lysine, polylysine, triolein, polyoxyethylene ethers and analogues thereof,
- 20 polidocanol alkyl ethers and analogues thereof, and mixtures thereof;
- d) mixing the mixed micellar composition resulting from steps a) to c) with a phenol selected from the group consisting of phenol, cresol and mixtures
- 25 thereof; and subsequently
- e) placing the formulation into an aerosol dispenser and charging the dispenser a propellant;
- wherein each of the absorption enhancing compounds are present in a concentration of from 1 to 10 wt./wt.%
- 30 of the total formulation, and the total concentration of alkali metal salicylate, alkali metal C8 to C22 alkyl

sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

The vigorous mixing may be accomplished using high speed stirrers, The vigorous mixing may be accomplished
5 by using high speed stirrers, e.g. magnetic stirrers or propellor stirrers, or by sonication.

In one embodiment, the mixed micellar formulation is formed by sonication of the aqueous micellar pharmaceutical agent composition in the presence of
10 lecithin.

Detailed Description of Preferred Embodiments

The present invention provides an improved method for delivery of macromolecular (high molecular weight) pharmaceutical agents, particularly through the
15 membranes in the nose, mouth, vagina or rectum. The preferred delivery is through oral and nasal cavities. The pharmaceutical agents cover a wide spectrum of agents, including proteins, peptides, hormones, vaccines and drugs. The molecular weights of the macromolecular
20 pharmaceutical agents are preferably above 1000, especially between 1000 and 2 000 000.

For example, hormones which may be administered with the present invention include thyroids, androgens, estrogens, prostaglandins, somatotropins, gonadotropins,
25 erythropoetin, interferons, interleukins, steroids and cytokins. Vaccines which may be administered with the present invention include bacterial and viral vaccines such as vaccines for hepatitis, influenza, tuberculosis, canary pox, chicken pox, measles, mumps, rubella,
30 pneumonia, BCG, HIV and AIDS. Bacterial toxoids which may be administered using the present invention include

diphtheria, tetanus, pseudomonas and mycobactrium tuberculosis. Examples of specific cardiovascular or thromobolytic agents include heparin, hirugen, hirulos and hirudin. Large molecules usefully administered with
5 the present invention include monoclonal antibodies, polyclonal antibodies and immunoglobins.

As will be understood, the concentration of the pharmaceutical agent is an amount sufficient to be effective in treating or preventing a disorder or to
10 regulate a physiological condition in an animal or human. The concentration or amount of pharmaceutical agent administered will depend on the parameters determined for the agent and the method of administration, e.g. oral, nasal. For example, nasal
15 formulations tend to require much lower concentrations of some ingredients in order to avoid irritation or burning of the nasal passages. It is sometimes desirable to dilute an oral formulation up to 10-100 times in order to provide a suitable nasal formulation.

20 The mixed micellar formulation is prepared by first preparing a first micellar composition which contains the pharmaceutically active agents, alkali metal C8 to C22 alkyl sulphate, edetate and alkali metal salicylate. For those compositions intended for administration
25 through the nasal, oral, vaginal or rectal cavities, the first micellar composition is then added to at least one of the absorption enhancing compounds to form a mixed micellar composition. At least one other absorption enhancing compound may also be added subsequently.

30 Preferably the first absorption enhancing compound is lecithin.

When making the aerosol formulation, the phenol and/or m-cresol and/or isotonic agent are then added. The formulation is then put into an aerosol dispenser and the dispenser charged with the propellant.

5 The preferred propellants are hydrogen-containing chlorofluorocarbons, hydrogen-containing fluorocarbons, dimethyl ether and diethyl ether. Even more preferred is hydrofluoroalkane (HFA) 134a (1,1,1,2 tetrafluoroethane).

10 Although the present invention has such wide applicability, the invention is described hereinafter with particular reference to insulin and its analogues, which are used for the treatment of diabetes.

As indicated hereinbefore, the compositions of the
15 present invention require that the pharmaceutical formulation be in mixed micellar form.

In the case of insulin, which is intended for administration through nasal or oral cavities, the first micellar solution may be made by adding a buffer
20 solution to powdered insulin, and then stirring until the powder is dissolved and a clear solution is obtained. A typical buffer solution is an aqueous solution of sodium salicylate and sodium lauryl sulphate and disodium edetate. Typical concentration of sodium
25 salicylate and sodium lauryl sulphate in the aqueous solution are about 3 to 20 wt./wt.% of each compound in the solution. Typically, insulin is present in the micellar solution in an amount which will give a concentration of about 2 to 4 wt./wt.% of the final
30 formulation. Typically the concentration may be about 10 wt./wt.% of the first micellar composition.

The micellar solution is then added slowly to the first absorption enhancing compound, e.g. lecithin while mixing vigorously, e.g. sonicating, to form a mixed micelle liposomal solution. At least one other
5 absorption enhancing compounds selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid,
10 linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein is then added. The mixing may be done with a high speed mixer or sonicator to ensure uniform micelle particle size distribution within
15 the formulation.

Each of the absorption enhancing compounds, when present, is in a concentration of from 1 to 10 wt./wt.% of the total formulation.

Preferred salts of hyaluronic acid are alkali metal
20 hyaluronates, alkaline earth hyaluronates and aluminium hyaluronate. The preferred salt is sodium hyaluronate. The preferred concentration of hyaluronic acid or pharmaceutically acceptable salts of hyaluronic acid is from 1 to 5 wt./wt.% of the total formulation. An even
25 more preferred range is from 1.5 to 3.5 wt./wt.% of the total formulation.

Other ingredients may be added to the mixed micellar solution. For example, flavouring agents, antioxidants, salts, protease inhibitors or other
30 pharmaceutically acceptable compound may be added.

In general the size of the micelle particles in the

solution is about 1 to 10 nm, and preferably from 1 to 5 nm. Such a size distribution ensures effective absorption of the formulation, and therefore the pharmaceutical agent, through the membranes, for example the membranes in the oral and nasal cavities.

The specific concentrations of the essential ingredients can be determined by relatively straightforward experimentation. For absorption through the nasal and oral cavities, it is often desirable to increase, e.g. double or triple, the dosage which is normally required through injection of administration through the gastrointestinal tract.

As will be understood, the amount of each component of the formulation will vary depending on the pharmaceutical agent and the site of application. Preferred formulations oral or nasal application have the following combinations: i) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H and sodium hyaluronate; ii) sodium lauryl sulphate, sodium salicylate, disodium edetate, lecithin and sodium hyaluronate; iii) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate and evening of primrose oil; iv) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H and bacitracin; v) sodium lauryl sulphate, sodium salicylate, disodium edetate, Phospholipon-H, sodium hyaluronate and bacitracin; and vi) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate, oleic acid and gamma linoleic acid.

For aerosol formulations, the addition of a mixture of phenol and m-cresol is preferred. Such an aerosol

formulation may then be charged to an aerosol dispenser and then charged with a propellant, preferably a non-CFC propellant.

The therapeutic compositions of the present invention may be stored at room temperature or at cold temperature. Storage of proteinic drugs is preferable at a cold temperature to prevent degradation of the drugs and to extend their shelf life.

As indicated hereinbefore, generally, oral and nasal are the favourite sites of the administration but the composition can be applied to the rectal and vaginal mucosa. According to the physiologically active peptide or protein used, the dosage form and the site of administration a specific administration method can be selected.

The composition of this invention is generally prepared as microfine mixed micellar particles (1 to 10 nm or less) by the virtue of its preparation methods used and suitable combinations of absorption enhancer characteristics.

For oral and nasal application, sprays are preferable, but also drops, chewable tablets, chewable gum and other suitable forms may be used. Utilization of atomizer or aerosol spray devices (metered dose inhalers or nebulizers) can be used to further reduce the particle size for effective inhalation from the nasal or oral cavity so the drug may successfully reach to the specific site and be absorbed. It is also possible to utilize a drug delivery system such that an enteric coating is applied to the gelatin capsule to cause the micelles to be released only in the duodenum

or in the proximity of the large intestine and not in the stomach.

The invention is illustrated by reference to the following examples.

5 Example 1

A first experiment was conducted to provide data for comparative purposes. This example does not fall within the scope of the present invention.

A solution was prepared using 0.5 g sodium lauryl
10 sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. To this solution 40 mg (1000 units) of insulin was added and dissolved completely while stirring, to give about 100 units/mL insulin solution.

15 In one set of tests, five healthy non-diabetic human volunteers were tested with insulin, by injection. In another set of tests the volunteers were tested with insulin, taken orally. The volunteers fasted from midnight prior to the test, with no food being taken
20 during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 100 units (1 mL volume per drop,
25 approximately 20 drops) of the above-prepared oral insulin (10 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

The average results for the five volunteers, of the
30 first day's trial (sub-cutaneous injection with 10 units) were as follows:

- 24 -

Table I

Time*:	0	15	30	60	75	90	120	150	180
Avg:	5.8	5.8	5.4	5.0	4.6	4.3	3.8	3.6	3.4
Time*:	210	240							
5 Avg:	4.2	4.5							

* time in minutes

The results for each of the five volunteers, of the second day's trial (oral drops with 100 units) were as follows:

10

Table II

Time*:	0	15	30	60	75	90	120	150	180
Subject Nos:									
1	6.2	5.8	5.2	5.0	4.9	5.0	5.0	4.8	4.7
2	5.8	5.4	5.0	4.7	4.9	4.3	5.0	5.5	5.2
15 3	4.8	4.6	4.3	4.3	4.4	4.6	4.8	4.7	5.2
4	6.6	6.1	5.8	5.5	5.1	4.9	5.0	5.0	5.9
5	6.0	5.8	5.7	5.5	5.1	4.8	4.7	4.9	5.0
Time*:		210	240						

Subject Nos:

20 1	5.5	6.0
2	5.8	6.1
3	5.5	5.1
4	6.2	6.8
5	5.9	6.7

25 * time in minutes

These tests indicate that compared to the injection method, oral insulin gives a faster onset of action and lowers blood glucose levels without creating hypoglycaemic condition. Due to the hepatic glucose production, there was a rebound effect. This is believed to be due to the incomplete absorption of insulin.

Example 2

Another experiment, not within the scope of the present invention, was performed for comparative purposes.

- 5 Oral insulin (100 units) was formulated in (Phospholipon-H, 10 mg) without any sodium lauryl sulphate, sodium salicylate, edetate or absorption enhancers, to evaluate its efficacy of blood glucose lowering in a fasted state, for healthy volunteers.
- 10 Volunteers were asked to fast overnight and not have any breakfast prior to dosing. Volunteers were asked to take this oral insulin formulation in their mouth and swallow it. Blood glucose levels were monitored every 15 minutes using Bayer's glucometer
- 15 Elite for 3 hours, and the average results for 5 volunteers are shown in Table III.

Table III

Time*:0	15	30	45	60	75	90	120	150	180
Avg: 5.6	5.8	5.8	5.7	5.7	5.8	5.7	5.7	5.8	5.7

20 * time in minutes

This indicates that orally administered insulin with lecithin alone has no effect on blood glucose lowering.

Example 3

- 25 A further experiment, not within the scope of the present invention, was performed for comparative purposes.

Oral insulin (100 units) was formulated with sodium salicylate and alkali metal edetate (both 5% by wt.) to

30 evaluate its efficacy of blood glucose lowering in fasted state in healthy volunteers.

Volunteers were asked to fast overnight and not have any breakfast prior to dosing. Volunteers were asked to take this oral insulin formulation in their mouth and swallow it. Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for 5 volunteers are shown in Table IV.

Table IV

Time*:	0	15	30	45	60	75	90	120	150	180
10 Avg:	5.8	5.8	5.8	5.9	5.8	5.9	5.7	5.9	6.2	6.0

* time in minutes

This indicates that orally administered insulin with sodium salicylate and alkali metal edetate alone has no effect on blood glucose lowering. In addition, 15 this formulation caused irritation and burning sensation, which lasted for several hours.

Example 4

A further experiment, not within the scope of the present invention, was performed for comparative 20 purposes.

Oral insulin (100 units) was formulated using sodium salicylate and alkali metal edetate (both 5% by wt.) with Phospholipon-H (10 mg) and tested on healthy subjects. Blood glucose levels were monitored every 15 25 minutes using Bayer's glucometer Elite for 3 hours and the results are shown in Table V.

Table V

Time*:	0	15	30	45	60	90	120	180
30 Avg:	5.3	5.3	5.3	5.4	5.6	5.7	5.7	5.8

* time in minutes

This indicates that orally administered insulin

with sodium salicylate, alkali metal edetate and Phospholipon-H has no effect on blood glucose lowering.

Example 5

Another experiment, not within the scope of the present invention, was performed for comparative purposes.

Oral insulin (50 units) was formulated using only alkali metal lauryl sulphate (5% by wt). Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for four volunteers are shown in Table VI.

Table VI

Time*:	0	15	30	60	90	120	180
Avg:	5.8	5.6	5.4	5.3	5.4	5.4	5.6

* time in minutes

This data shows that orally administered insulin with only alkali metal lauryl sulphate has little metabolic effect on the blood glucose lowering in healthy subjects. This formulation caused substantial burning sensation and irritation in the subjects and lasted for two days.

Example 6

Yet another experiment, within the scope of the present invention, was performed.

Mixed micellar oral insulin (50 units) was formulated using alkali metal lauryl sulphate and sodium salicylate (both 4.4% by wt.) and alkali metal edetate (2.2% by wt) with Phospholipon-H (10 mg) and tested on healthy volunteers.

The method involved mixing the sodium lauryl sulphate, sodium salicylate and alkali metal edetate

with water in a beaker with a magnetic stirrer at medium speed until the ingredients were dissolved, to form buffer solution. Insulin powder was placed in a beaker and to this powder was added the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution obtained. The micellar solution so formed was stored in clean glass bottles and refrigerated.

10 Mixed micellar liposomal insulin was then prepared in a glass beaker, in which was placed the Phospholipon-H and a small amount of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10 minutes to ensure complete dissolution of the
15 Phospholipon-H. To this solution was added the micellar insulin solution very slowly, drop wise, using glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30 minutes at a high speed to ensure uniform micellar particle size
20 distribution.

Samples of the mixed micellar solution were taken orally by the volunteers.

Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and
25 the average results for 5 volunteers are shown in Table VII.

Table VII

Time*:	0	15	30	45	60	90	120	150	180
Avg:	6.5	6.1	5.5	5.3	5.3	5.4	5.5	5.5	5.5

30 * time in minutes

This data shows that orally administered insulin

with alkali metal lauryl sulphate combined with the sodium salicylate and alkali metal edetate with Phospholipon-H has a small metabolic effect on blood glucose levels in healthy volunteers.

5 Example 7

An experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

Oral insulin (50 units) was formulated using alkali
10 metal lauryl sulphate and sodium salicylate (both 4.4% by wt.) and alkali metal edetate (2.2% by wt.) with Phospholipon-H (10 mg) and sodium hyaluronate (1.1% by wt). This formulation was tested on healthy subjects under fasting condition.

15 The method involved mixing the sodium lauryl sulphate, sodium salicylate and alkali metal edetate with water in a beaker with a magnetic stirrer at medium speed until the ingredients were dissolved, to form buffer solution. Insulin powder was placed in a beaker
20 and to this powder was added the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution obtained. The micellar solution so formed was stored in clean glass bottles and
25 refrigerated.

Mixed micellar liposomal insulin was then prepared in a glass beaker, in which was placed the Phospholipon-H and a small amount of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10
30 minutes to ensure complete dissolution of the Phospholipon-H. To this solution was added the micellar

insulin solution very slowly, drop wise, using glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30 minutes at a high speed to ensure uniform micellar particle size distribution. The hyaluronate and small amounts of menthol and sorbitol were then added, with continuous stirring.

Samples of the mixed micellar solution were taken orally by the volunteers.

10 Blood glucose levels were monitored every 15 minutes using Bayer's glucometer Elite for 3 hours and the average results for 5 volunteers are shown in Table VIII.

Table VIII

15	Time*:	0	15	30	45	60	90	120	150	180
	Avg:	6.5	5.9	5.6	5.4	4.9	5.0	4.9	5.2	5.4

* time in minutes

This data shows that orally administered insulin with alkali metal lauryl sulphate, sodium salicylate, 20 alkali metal edetate, Phospholipon-H and sodium hyaluronate has resulted in lowering of blood glucose levels in healthy subjects better than the above mentioned formulations.

Example 8

25 A further experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g 30 disodium edetate dissolved in 10 mL of water. The solution was added to insulin and mixed, to form

micellar insulin.

Separately, 100 mg of powdered Phosphatidylcholin-H was added to a glass beaker and to this powder was added 10 mL 50% ethanol. The powder was dissolved completely.

5 To this solution 16 mg (400 units) of micellar insulin solution dissolved in 3 mL of the buffer solution to (give 30 units/mL insulin solution) was added slowly with vigorous mixing, to form a mixed micellar solution. To this was added 0.6 mL of sodium hyaluronate and 0.2

10 mL of 2% menthol solution containing 3% sorbitol.

In one set of tests, ten Type II diabetic human volunteers who took insulin, by injection three times a day, were studied. In another set of tests the volunteers were tested with insulin, taken orally. The

15 volunteers fasted from midnight prior to the test, with no food being taken during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the

20 volunteers received 30 units (1 mL volume per drop, approximately 20 drops) of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

25 The results, showing the average for the ten volunteers, were as shown on the following page:

Table IX

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose (30 units)	Injection (10 units)
5		
0	6.4	6.8
15	5.8	6.9
30	5.4	6.1
45	5.3	5.8
10 60	5.3	5.8
75	5.2	5.8
90	5.2	5.4
105	5.2	5.4
120	5.1	5.2
15 135	5.1	5.1
150	5.2	4.9
165	5.3	4.9
180	5.3	4.8
195	5.4	4.8
20 210	5.4	5.2
225	5.6	5.2
240	5.6	5.4

The results show that the oral insulin formulation of the present invention, at a dosage of three times higher than the injected level, is comparable to the injected insulin.

Example 9

This example illustrates a method for making a mixed micellar formulation according to the present invention.

In a 250 mL capacity glass beaker was added 5 g

sodium lauryl sulphate, 5 g sodium salicylate and 2.5 g edetate. The beaker was placed on the hot plate with a magnetic stirrer. To this dry powder mixture was added 100 mL distilled water and the mixture was stirred, 5 using the magnetic stir bar, at a medium speed until all the powder was dissolved. The buffer solution was stored in a clean glass bottle at room temperature (pH 6.5).

A micellar insulin solution was then prepared in a 10 50 mL capacity glass beaker, into which was placed 11.54 mg insulin powder. To this powder was added 10 mL of the buffer solution. The solution was continuously stirred using a magnetic stir bar until all of the insulin powder was dissolved and a clear solution 15 obtained. The micellar solution so formed was stored in clean glass bottles and refrigerated.

A 2% menthol solution was then prepared from 100 mg menthol crystals, dissolved in 5 mL ethanol. To this solution was added 5 mg FD & C blue dye. The solution 20 was stirred for 10 minutes and stored in a glass bottle at room temperature.

Mixed micellar liposomal insulin was then prepared in a 50 mL glass beaker, in which was placed 100 mg of phosphatidylcholine (Sigma, type I=EH, hydrogenated). To 25 this powder was added 10 mL of isopropyl alcohol. The mixture was stirred at a high speed (1000 rpm) for about 10 minutes to ensure complete dissolution of the phosphatidylcholine. To this solution was added the micellar insulin solution very slowly, drop wise, using 30 glass dropper, with continuous stirring at a high speed. The solution was stirred continuously for another 30

minutes at a high speed to ensure uniform micellar particle size distribution. To this solution was added 1 mL of the 2% menthol solution and 50 mg sodium hyaluronate. The semi-clear, translucent, light blue colour, liposomal insulin mixed micellar solution (final volume 15 mL) was stored in a clean glass bottle and refrigerated. The solution had a pH of 6.5.

If the phosphatidylcholine powder does not dissolve completely, then heating up to about 45°C may be required, e.g. using a water bath.

It has been found that if the micellar insulin composition is not added slowly, then the mixed micellar formulation will not be formed and the formulation will be gelatinous and sticky.

15 Example 10

The formulation of Example 9 was tested in a manner similar to that indicated in Example 8 except that the formulation of the present invention was administered nasally.

20 On the first day, the ten volunteers each received 10 units insulin injection (regular fast acting, Eli Lilly). On the second day, the volunteers received 20 units of the "oral" insulin of Example 9 (2 times the injection dose). The "oral" insulin was administered as 25 drops (0.4 mL volume per drop, approximately 4 large drops in total, i.e. two drops in each nostril).

The results, showing the average for the ten volunteers, are shown on the following page:

Table X

		Blood glucose levels (mmol/L)	
	Time (minutes)	Nasal Dose	Injection
		(20 units)	(10 units)
5	0	7.4	6.8
	15	6.7	7.0
	30	5.9	6.8
	45	5.3	6.3
	60	5.0	6.3
10	75	5.2	5.8
	90	5.1	5.2
	105	5.0	5.0

Table X (continued)

		Blood glucose levels (mmol/L)	
	Time (minutes)	Nasal Dose	Injection
		(20 units)	(10 units)
15	120	4.6	5.2
	135	4.5	4.2
	150	4.3	4.6
	165	4.3	4.0
	180	4.8	4.1
20	195	5.3	4.3
	210	5.4	4.5
	225	5.7	4.7
	240	5.6	5.0

The results show that the nasal insulin formulation of the present invention, at a dosage of twice the injected level, is comparable to the injected insulin.

Example 11

30 The formula of Example 9 was taken and tests performed to determine the insulin action on meal

glucose on healthy volunteers.

Usually, diabetic patients take an insulin injection 30 minutes prior to a meal, because injected insulin takes a long time to take effect. Injected
5 insulin is slowly absorbed into bloodstream within 60 minutes and has metabolic effect on meal glucose levels.

The mixed micellar formulation of Example 9 was tested in healthy volunteers under controlled conditions to determine the oral insulin effect on meal glucose
10 when compared to injected insulin.

In one set of tests, ten healthy non-diabetic human volunteers were tested with insulin, by injection. In another set of tests the volunteers were tested with insulin, taken orally. The volunteers fasted from
15 midnight prior to the tests, with food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units
20 insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 30 units of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by
25 Bayer's Glucometer Elite. The results are shown on the following page:

Table XI

Blood glucose levels (mmol/L)		
Time (minutes)	Oral Dose (30 units)	Injection (10 units)
5		
0	5.7	5.5
15	5.2	5.6
30	5.0	5.4
45	5.3	5.4
10 60	5.4	5.6
75	6.3	6.6
90	6.9	7.0
105	6.0	5.9
120	5.8	5.6
15 135	5.5	5.1
150	5.1	4.8
165	4.9	4.6
180	4.8	4.3

The results indicate that the oral insulin helps
 20 control meal glucose levels in healthy volunteers when
 compared to injected insulin.

Example 12

The mixed micellar formulation of Example 9 was
 tested in diabetic volunteers under controlled
 25 conditions to determine the oral insulin effect on meal
 glucose when compared to injected insulin.

In one set of tests, ten Type II diabetic human
 volunteers who took insulin, by injection three times a
 day, were studied. In another set of tests the
 30 volunteers were tested with insulin, taken orally. The
 volunteers fasted from midnight prior to the tests, with

food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units 5 insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the volunteers received 30 units of the above-prepared oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by 10 Bayer's Glucometer Elite.

The average results for the 10 volunteers were as follows:

Table XII

		Blood glucose levels (mmol/L)	
15	Time (minutes)	Oral Dose	Injection
		(30 units)	(10 units)
	0	8.8	8.7
	15	8.1	8.8
	30	8.0	8.9
20	45	8.4	10.1
	60	10.2	11.8
	75	11.8	11.8
	90	12.3	12.2
	105	10.8	11.2
25	120	9.6	10.4
	135	8.1	8.4
	150	6.9	7.3
	165	6.2	6.5
	180	4.8	4.3

30 The results indicate that oral insulin helps to control meal glucose levels in diabetic patients when

compared to injected insulin.

Example 13

A chewable gum insulin formulation was prepared by vigorously stirring the liposomal insulin mixed micellar solution of Example 9 while adding guar gum, beeswax, powdered acacia, oleic acid, gamma-linoleic acid and sorbitol. For each 30 units of insulin, the mixture contained 100 mg guar gum, 50 mg beeswax, 50 mg powdered acacia, 100 mg oleic acid, 100 mg gamma-linoleic acid and 1 mL 3% sorbitol in ethanol solution. The mixture was then poured into a flat tray coated with polytetrafluoroethylene until the mixture was about 10 mm deep. The mixture then solidified and after solidification was cut into sticks about 1 cm by 3 cm. Each stick contained about 30 units insulin.

The mixed micellar formulation in chewable stick form was tested in diabetic volunteers under controlled conditions to determine the oral insulin effect on meal glucose when compared to injected insulin.

In one set of tests, five Type II diabetic human volunteers who took insulin, by injection three times a day, were studied. In another set of tests the volunteers were tested with the chewable gum insulin, taken orally. The volunteers fasted from midnight prior to the tests, with food being taken 30 minutes after dosing. The meals were standard Sastacal 240 mL liquid diet approved by the Diabetic Society, containing 400 calories.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, available from Eli Lilly). On the second day, the

volunteers received 30 units of the above-prepared chewable gum oral insulin (3 times the injection dose). In both tests, blood glucose levels were monitored every 15 minutes by Bayer's Glucometer Elite.

- 5 The average results for the five volunteers were as follows:

Table XIII

		Blood glucose levels (mmol/L)	
Time (minutes)		Oral Dose	Injection
		(30 units)	(10 units)
10	0	9.1	8.8
	15	9.3	8.2
	30	9.3	8.0
	45	10.2	8.4
15	60	11.2	9.2
	75	12.1	10.3
	90	12.9	11.8
	105	13.2	11.6
	120	12.8	11.0
20	135	12.2	10.2
	150	11.6	9.6
	165	11.0	9.5
	180	10.6	9.1
	195	10.0	8.7
25	210	9.5	8.2
	225	8.8	8.0
	240	8.2	7.5

Example 14

Another experiment, within the scope of the present invention, was performed. In this example, the formulation was for oral administration.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. The solution was added to 8 mg (200 units) insulin and 5 mixed, to form micellar insulin.

To this micellar solution were added 0.2 g bacitracin and 0.5 g evening of primrose oil and the solution was mixed vigorously to form a mixed micellar insulin solution (about 20 units/mL).

10 Six human volunteers were studied. The volunteers fasted from midnight prior to the test, with no food being taken during the 4 hour study.

On the first day, the volunteers received 10 units insulin by injection (regular fast acting insulin, 15 available from Eli Lilly). On the second day, the volunteers received 20 units of the above-prepared oral insulin (twice the injection dose). In both tests, blood glucose levels were monitored at intervals by Bayer's Glucometer Elite.

20 The results, showing the average for the six volunteers, were as follows:

Table XIV

Time (minutes)	Blood glucose levels (mmol/L)	
	Oral Dose (20 units)	Injection (10 units)
25 0	8.8	7.9
15	8.4	7.9
30	8.1	8.2
45	7.4	8.3
30 60	6.3	7.6
90	5.1	6.2

120	5.0	5.2
150	4.8	4.6
180	5.1	3.9
210	5.3	4.4
5 240	5.6	5.2

The results show that the oral insulin formulation of the present invention, at a dosage of twice the injected level, is comparable to the injected insulin.

Example 15

10 A further experiment was performed to show another method of making the mixed micellar formulation of the present invention.

 In a 250 mL round bottom flask was added 100-mg of saturated lecithin powder (Phospholipon-90H) purchased
15 from the American Lecithin Co. To this powder was added 5 mL of absolute ethanol (USP grade). The flask was then attached to a rotary evaporator equipped with the vacuum pump and nitrogen inlet for inert atmosphere condition to minimize oxidation of the lecithin. The
20 flask was rotated at 100-150 rpm under vacuum. The solution in the flask was heated to 60°C by means of water bath to dissolve the powder completely. After complete dissolution of the powder, heating was stopped and the rotation speed was increased to 300 rpm, under
25 vacuum in nitrogen atmosphere until the alcohol evaporated completely, leaving a uniform film on the side of the flask. The rotation was continued for at least 30 minutes to ensure uniform coating of film on the wall and complete solvent removal. After 30
30 minutes the rotation was stopped and the vacuum was released.

To this flask was added micellar insulin solution which had been prepared from an aqueous solution of insulin, sodium lauryl sulphate, sodium salicylate and disodium edetate. The flask was shaken with the help of shaker plate. Shaking was continued for at least 30 minutes and then the solution was sonicated with a high frequency sonicating probe for another 60 minutes in order to form small uniform mixed micelles. The mixed micelles so obtained were analyzed by Malvern Zeta (trade mark) particle size distribution measurement equipment equipped with the laser light scattering device. The mixed micelles particle size distribution obtained by this method was between 2 and 9nm. To this solution was added 1 mL of 2% menthol solution and 50 mg sodium hyaluronate. The semi-clear, translucent, light blue colour solution (final volume 10 mL) was stored in a clean glass bottle and refrigerated. The solution had a pH of 6.5.

Example 16

Another experiment, within the scope of the present invention, was performed.

A buffer solution was prepared using 0.5 g sodium lauryl sulphate, 0.5 g sodium salicylate and 0.25 g disodium edetate dissolved in 10 mL of water. The solution was added to 8 mg (200 units) insulin and mixed, to form micellar insulin.

To this micellar solution were added 0.5 g borage oil and the solution was mixed vigorously to form a mixed micellar insulin solution (about 20 units/mL).

CLAIMS:

1. A process for making a pharmaceutical composition suitable for delivery through mucosal membranes comprising:
 - 5 a) preparing a pharmaceutical agent composition in micellar form in an aqueous medium which has an alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition, an alkali metal C8 to C22 alkyl sulphate in
10 a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition and a pharmaceutically acceptable edetate in a concentration of from 1 to 10 wt./wt.% of the aqueous micellar pharmaceutical agent composition;
15 b) slowly adding the micellar proteinic pharmaceutical agent composition, while mixing, to at least one absorption enhancing compound, while continuing to mix vigorously, said absorption enhancing compounds being selected from the group consisting of lecithin,
20 hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, menthol, trihydroxy oxo cholanylglycine
25 and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing compound is present in a concentration of from 1 to
30 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal

C8 to C22 alkyl sulphate, edetate and absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

2. A process according to Claim 1 wherein there is an additional step of adding, while continuing mixing, at least one absorption enhancing compound different to that added in step b), selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin, lysine, polylysine, triolein and mixtures thereof.

3. A process according to Claim 1 wherein the absorption enhancing compound in step b) is selected from the group consisting of saturated phospholipid, unsaturated phospholipid, phosphatidylcholine, phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, lecithin, lysolecithin and mixtures thereof.

4. A process according to Claim 1 wherein one of the absorption enhancing compounds is lecithin and another absorption enhancing compound is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid and mixtures thereof, the concentration such absorption enhancing compound being from about 1 to about 5 wt./wt.%.

5. A process according to Claim 1 wherein the micellar absorption enhancing compounds comprise combinations selected from the group consisting of i) saturated

phospholipid and sodium hyaluronate, ii) saturated phospholipid and glycolic acid, iii) lecithin and sodium hyaluronate and iv) saturated phospholipid, glycolic acid and lactic acid.

- 5 6. A process according to Claim 1 wherein the proteinic pharmaceutical agent is selected from the group consisting of insulin, heparin, so-called low molecular weight heparin, hirulog, hirugen, huridin, interferons, interleukins, cytokines, mono and
- 10 polyclonal antibodies, chemotherapeutic agents, vaccines, glycoproteins, bacterial toxoids, hormones, calcitonins, insulin like growth factors (IGF), glucagon like peptides (GLP-1), large molecule antibiotics, protein based thrombolytic compounds, platelet
- 15 inhibitors, DNA, RNA, gene therapeutics, antisense oligonucleotides, opioids, narcotics, analgesics, NSAIDS, steroids, hypnotics, pain killers and morphine.

7. A process according to Claim 1 wherein in step b) the micellar proteinic pharmaceutical agent composition
- 20 is added to lecithin, with sonication, to form a mixed micellar composition; and

- c) while continuing to mix, adding at least one absorption enhancing compound selected from the group consisting of hyaluronic acid, pharmaceutically
- 25 acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of primrose oil, trihydroxy oxo cholanylglycine, glycerin, polyglycerin,
- 30 lysine, polylysine, triolein and mixtures thereof;

wherein the amount of lecithin and the absorption

enhancing compound are each present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of alkali metal salicylate, alkali metal C8 to C22 alkyl sulphate, edetate and
5 absorption enhancing compounds is less than 50 wt./wt.% of the formulation.

8. A process according to Claim 1 wherein the absorption enhancing compound is formed into a film prior to the addition of the micellar pharmaceutical
10 agent composition.

9. A process according Claim 1 wherein subsequent to the addition of the micellar pharmaceutical agent composition a second absorption enhancing compound is added, said second absorption enhancing compound being
15 different from the absorption enhancing compound previously used.

10. A process according to Claim 1 a phenol selected from the group consisting of phenol, methyl phenol and mixtures thereof are added to the micellar formulation
20 and the resulting formulation placed in a container, and the container is subsequently charged with a propellant.

11. A process according to Claim 10 wherein the propellant is selected from the group consisting of tetrafluoroethane, tetrafluoropropane,
25 dimethylfluoropropane, heptafluoropropane, dimethyl ether, n-butane and isobutane.

12. A process according to Claim 1 wherein the pharmaceutical agent is insulin.

13. A process according to Claim 11 wherein the
30 pharmaceutical agent is insulin.

14. A mixed micellar pharmaceutical formulation

comprising a pharmaceutical agent in micellar form, water, an alkali metal C8 to C22 alkyl sulphate in a concentration of from 1 to 10 wt./wt.% of the total formulation, a pharmaceutically acceptable edetate in a
5 concentration of from 1 to 10 wt./wt.% of the total formulation, at least one alkali metal salicylate in a concentration of from 1 to 10 wt./wt.% of the total formulation, and at least one absorption forming compound, said absorption forming compounds being
10 selected from the group consisting of lecithin, hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid, octylphenoxypolyethoxyethanol, glycolic acid, lactic acid, chamomile extract, cucumber extract, oleic acid, linolenic acid, borage oil, evening of
15 primrose oil, menthol, trihydroxy oxo cholanylglycine and pharmaceutically acceptable salts thereof, glycerin, polyglycerin, lysine, polylysine, polidocanol alkyl ethers and analogues thereof, triolein and mixtures thereof, wherein the amount of each absorption enhancing
20 compound is present in a concentration of from 1 to 10 wt./wt.% of the total formulation, and the total concentration of absorption enhancing compounds are less than 50 wt./wt.% of the formulation.

15. A mixed micellar pharmaceutical formulation
25 according to Claim 14, in which one of the absorption enhancing compounds is lecithin.

16. A formulation according to Claim 14 wherein the alkali metal C8 to C22 alkyl sulphate is sodium lauryl sulphate and the alkali metal salicylate is sodium
30 salicylate.

17. A formulation according to Claim 15 wherein the

lecithin is selected from the group consisting of saturated phospholipid, unsaturated phospholipid, phosphatidylcholine, phosphatidyl serine, sphingomyelin, phosphatidylethanolamine, cephalin, lysolecithin and

5 mixtures thereof.

18. A formulation according to Claim 15 wherein the other absorption enhancing compound is selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid and

10 mixtures thereof, the concentration such absorption enhancing compound being from about 1 to about 5 wt./wt. %.

19. A formulation according to Claim 14 wherein the formulation comprises combinations selected from the group consisting of i) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated phospholipid and sodium hyaluronate; ii) sodium lauryl sulphate, sodium salicylate, disodium edetate, lecithin and sodium hyaluronate; iii) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate and evening of primrose oil; iv) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated phospholipid and bacitracin; v) sodium lauryl sulphate, sodium salicylate, disodium edetate, saturated phospholipid, sodium hyaluronate and bacitracin; and vi) sodium lauryl sulphate, sodium salicylate, disodium edetate, sodium hyaluronate, oleic acid and gamma linoleic acid.

20. A formulation according to Claim 14 wherein the pharmaceutical agent is selected from the group consisting of insulin, heparin, so-called low molecular

weight heparin, hirulog, hirugen, huridine, interferons, interleukins, cytokins, mono and polyclonal antibodies, chemotherapeutic agents, vaccines, glycoproteins, bacterial toxoids, hormones, calcitonins, insulin like
5 growth factors (IGF), glucagon like peptides (GLP-1), large molecule antibiotics, protein based thrombolytic compounds, platelet inhibitors, DNA, RNA, gene therapeutics antisense oligonucleotides, opioids, narcotics, analgesics, NSAIDS, steroids, hypnotics, pain
10 killers and morphine.

21. A formulation according to Claim 14 wherein the pharmaceutical agent is insulin.

22. A formulation according to Claim 21 in which the absorption enhancing compounds are lecithin and a second
15 absorption enhancing compound selected from the group consisting of hyaluronic acid, pharmaceutically acceptable salts of hyaluronic acid and mixtures thereof.

23. A formulation according to Claim 14 wherein the
20 formulation additionally comprises a phenol selected from the group consisting of phenol, methyl phenol and mixtures thereof.

24. A formulation according to Claim 23 wherein the formulation is contained in an aerosol container and the
25 container is charged with a propellant.

25. A formulation according to Claim 24 wherein the propellant is selected from the group consisting of tetrafluoroethane, tetrafluoropropane, dimethylfluoropropane, heptafluoropropane, dimethyl
30 ether, n-butane and isobutane.

INTERNATIONAL SEARCH REPORT

In: ational Application No

PCT/CA 99/00106

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K38/28 A61K9/107

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 36352 A (MODI PANKAJ ;CHANDARANA SUBASH (CA)) 21 November 1996 see page 3, line 2-27 see page 5, line 7-15 see page 5, line 35 - page 6, line 25 see page 7, line 4-5 see claims ----	1-7, 9, 12-22
A	WO 96 19197 A (ASTRA AB ;BAECKSTROEM KJELL (SE); DAHLBAECK MAGNUS (SE); JOHANSSON) 27 June 1996 see page 2, line 15-24 see page 3, line 26-30 see page 4, line 10-27 see page 5, line 22 - page 6, line 14 see examples see claims 1,8-12,15-17,24-26,43 ----- -/--	1,6, 10-14, 24,25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 June 1999

Date of mailing of the international search report

10/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

La Gaetana, R

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/CA 99/00106

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	YAMAMOTO A ET AL: "Pulmonary absorption enhancement of peptides by absorption enhancers and protease inhibitors" JOURNAL OF CONTROLLED RELEASE, vol. 41, no. 1, 1 August 1996, page 57-67 XP004037571 see abstract see pages 60-62, "Results", paragraphs 3.1-3.3 see page 64, column 2, line 24-page 65, column 1, line 5	1,14
A	WO 94 26302 A (GENENTECH INC) 24 November 1994 see page 3, line 2-22 see page 6, line 18-27 see page 7, line 4-23 see claims 1,11,12,14	1,10,11, 14,23,24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 99/00106

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9636352 A	21-11-1996	US 5653987 A	05-08-1997
		AU 5642396 A	29-11-1996
		CA 2210996 A	21-11-1996
		EP 0813421 A	29-12-1997
WO 9619197 A	27-06-1996	AU 702879 B	11-03-1999
		AU 4359196 A	10-07-1996
		BR 9510501 A	13-01-1998
		CA 2206736 A	27-06-1996
		CN 1171046 A	21-01-1998
		CZ 9701945 A	15-10-1997
		EP 0797431 A	01-10-1997
		FI 972657 A	19-06-1997
		HU 77701 A	28-07-1998
		JP 10510827 T	20-10-1998
		NO 972781 A	16-06-1997
		PL 320824 A	10-11-1997
		SK 81397 A	05-11-1997
		ZA 9510752 A	24-06-1996
WO 9426302 A	24-11-1994	AT 173632 T	15-12-1998
		AU 683469 B	13-11-1997
		AU 6782294 A	12-12-1994
		CA 2159602 A	24-11-1994
		CN 1122575 A	15-05-1996
		CZ 9502955 A	17-04-1996
		DE 69414841 D	07-01-1999
		DE 69414841 T	02-06-1999
		EP 0697887 A	28-02-1996
		ES 2126114 T	16-03-1999
		FI 955136 A	27-10-1995
		HU 72980 A	28-06-1996
		JP 8510242 T	29-10-1996
		NO 954544 A	10-11-1995
		NZ 266354 A	24-06-1997
		PL 311639 A	04-03-1996
		SK 137895 A	05-02-1997
		ZA 9402955 A	30-10-1995